



United States
Department of
Agriculture

NRCS

Natural
Resources
Conservation
Service

A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for Blair County, Pennsylvania, and Huntingdon County, Pennsylvania

The Bookhammer Rural Lifestyle Farm



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<http://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means

for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.


Contents

Preface	2
Soil Map	5
Soil Map.....	6
Legend.....	7
Map Unit Legend.....	9
Map Unit Descriptions.....	9
Blair County, Pennsylvania.....	12
BrB—Brinkerton silt loam, 3 to 8 percent slopes.....	12
BuB—Buchanan gravelly silt loam, 3 to 8 percent slopes.....	13
BxB—Buchanan extremely stony silt loam, 3 to 8 percent slopes.....	15
CbB—Clarksburg silt loam, 3 to 8 percent slopes.....	16
EmB—Edom-Weikert complex, 3 to 8 percent slopes.....	17
MuB—Murrill gravelly silt loam, 3 to 8 percent slopes.....	19
MuD—Murrill gravelly silt loam, 15 to 25 percent slopes.....	20
MxB—Murrill extremely stony silt loam, 3 to 8 percent slopes.....	21
OuC—Opequon silty clay loam, 8 to 15 percent slopes.....	22
OuD—Opequon silty clay loam, 15 to 25 percent slopes.....	23
UD—Udfluvents-Dystrochrepts complex.....	25
Huntingdon County, Pennsylvania.....	28
Ba—Barbour soils.....	28
HeD—Hagerstown-Rock outcrop complex, 5 to 25 percent slopes.....	29
OpD3—Opequon clay loam, 15 to 25 percent slopes, eroded.....	30
Soil Information for All Uses	32
Soil Reports.....	32
Land Classifications.....	32
Hydric Soils.....	32
Vegetative Productivity.....	35
Nonirrigated Yields by Map Unit.....	35

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map



Map Scale: 1:6,070 if printed on A landscape (11" x 8.5") sheet.


Meters




Map projection: Web Mercator Corner coordinates: WGS84 Edge tics: UTM Zone 17N WGS84


MAP LEGEND


Area of Interest (AOI)

 Area of Interest (AOI)




















Soils






 Soil Map Unit Polygons

 Soil Map Unit Lines


 Soil Map Unit Points

Special Point Features






-  Blowout
-  Borrow Pit
-  Clay Spot
-  Closed Depression
-  Gravel Pit
-  Gravelly Spot
-  Landfill
-  Lava Flow
-  Marsh or swamp
-  Mine or Quarry
-  Miscellaneous Water
-  Perennial Water
-  Rock Outcrop
-  Saline Spot
-  Sandy Spot
-  Severely Eroded Spot
-  Sinkhole
-  Slide or Slip
-  Sodic Spot

-  Spoil Area
-  Stony Spot
-  Very Stony Spot
-  Wet Spot
-  Other
-  Special Line Features


Water Features

 Streams and Canals

Transportation

-  Rails
-  Interstate Highways
-  US Routes
-  Major Roads
-  Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:20,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL: <http://websoilsurvey.nrcs.usda.gov>
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Blair County, Pennsylvania
 Survey Area Data: Version 9, Nov 16, 2015

Soil Survey Area: Huntingdon County, Pennsylvania
 Survey Area Data: Version 8, Nov 16, 2015

Your area of interest (AOI) includes more than one soil survey area. These survey areas may have been mapped at different scales, with a different land use in mind, at different times, or at different levels of detail. This may result in map unit symbols, soil properties, and interpretations that do not completely agree across soil survey area boundaries.

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Custom Soil Resource Report

MAP LEGEND

MAP INFORMATION

Date(s) aerial images were photographed: Oct 6, 2011—Oct 17, 2011

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Blair County, Pennsylvania (PA013)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
BrB	Brinkerton silt loam, 3 to 8 percent slopes	11.9	13.4%
BuB	Buchanan gravelly silt loam, 3 to 8 percent slopes	5.9	6.7%
BxB	Buchanan extremely stony silt loam, 3 to 8 percent slopes	13.3	14.9%
CbB	Clarksburg silt loam, 3 to 8 percent slopes	9.9	11.1%
EmB	Edom-Weikert complex, 3 to 8 percent slopes	0.1	0.1%
MuB	Murrill gravelly silt loam, 3 to 8 percent slopes	3.5	3.9%
MuD	Murrill gravelly silt loam, 15 to 25 percent slopes	7.9	8.9%
MxB	Murrill extremely stony silt loam, 3 to 8 percent slopes	8.4	9.5%
OuC	Opequon silty clay loam, 8 to 15 percent slopes	7.7	8.7%
OuD	Opequon silty clay loam, 15 to 25 percent slopes	11.3	12.7%
UD	Udifluvents-Dystrochrepts complex	7.5	8.4%
Subtotals for Soil Survey Area		87.4	98.3%
Totals for Area of Interest		88.9	100.0%

Huntingdon County, Pennsylvania (PA061)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
Ba	Barbour soils	1.2	1.4%
HeD	Hagerstown-Rock outcrop complex, 5 to 25 percent slopes	0.1	0.1%
OpD3	Opequon clay loam, 15 to 25 percent slopes, eroded	0.2	0.2%
Subtotals for Soil Survey Area		1.5	1.7%
Totals for Area of Interest		88.9	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

Custom Soil Resource Report

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

Custom Soil Resource Report

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Blair County, Pennsylvania

BrB—Brinkerton silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 16bh
Elevation: 300 to 3,000 feet
Mean annual precipitation: 30 to 65 inches
Mean annual air temperature: 46 to 59 degrees F
Frost-free period: 120 to 217 days
Farmland classification: Not prime farmland

Map Unit Composition

Brinkerton and similar soils: 75 percent
Minor components: 25 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Brinkerton

Setting

Landform: Depressions
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Base slope
Down-slope shape: Concave
Across-slope shape: Concave
Parent material: Local fine-silty colluvium derived from sedimentary rock

Typical profile

H1 - 0 to 9 inches: silt loam
H2 - 9 to 18 inches: silty clay loam
H3 - 18 to 46 inches: silty clay loam
H4 - 46 to 65 inches: channery silt loam

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 15 to 34 inches to fragipan
Natural drainage class: Poorly drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: About 0 to 6 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 3.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: D
Hydric soil rating: Yes

Minor Components

Ernest

Percent of map unit: 10 percent
Hydric soil rating: No

Custom Soil Resource Report

Berks

Percent of map unit: 5 percent
Landform: Ridges, valleys
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex, linear
Across-slope shape: Convex, linear
Hydric soil rating: No

Laidig

Percent of map unit: 5 percent
Landform: Mountains
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Lower third of mountainflank
Down-slope shape: Concave
Across-slope shape: Concave
Hydric soil rating: No

Atkins

Percent of map unit: 3 percent
Landform: Flood plains
Down-slope shape: Concave
Across-slope shape: Concave
Hydric soil rating: Yes

Philo

Percent of map unit: 2 percent
Hydric soil rating: No

BuB—Buchanan gravelly silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 16bk
Elevation: 300 to 3,000 feet
Mean annual precipitation: 35 to 55 inches
Mean annual air temperature: 45 to 59 degrees F
Frost-free period: 120 to 217 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Buchanan and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Buchanan

Setting

Landform: Mountain slopes, valley sides
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Lower third of mountainflank, base slope

Custom Soil Resource Report

Down-slope shape: Concave, linear

Across-slope shape: Linear, concave

Parent material: Mountain slope colluvium derived from sedimentary rock

Typical profile

H1 - 0 to 8 inches: gravelly loam

H2 - 8 to 32 inches: gravelly loam

H3 - 32 to 65 inches: gravelly loam

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 20 to 36 inches to fragipan

Natural drainage class: Moderately well drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 18 to 36 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Low (about 4.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: C

Hydric soil rating: No

Minor Components

Andover

Percent of map unit: 6 percent

Landform: Depressions

Landform position (three-dimensional): Mountainbase

Down-slope shape: Concave

Across-slope shape: Concave

Hydric soil rating: Yes

Berks

Percent of map unit: 3 percent

Hydric soil rating: No

Bedington

Percent of map unit: 3 percent

Hydric soil rating: No

Philo

Percent of map unit: 3 percent

Hydric soil rating: No

BxB—Buchanan extremely stony silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 16bm
Elevation: 300 to 3,000 feet
Mean annual precipitation: 35 to 55 inches
Mean annual air temperature: 46 to 59 degrees F
Frost-free period: 120 to 217 days
Farmland classification: Not prime farmland

Map Unit Composition

Buchanan and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Buchanan

Setting

Landform: Mountain slopes, valley sides
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Lower third of mountainflank, base slope
Down-slope shape: Concave, linear
Across-slope shape: Linear, concave
Parent material: Mountain slope colluvium derived from sedimentary rock

Typical profile

H1 - 0 to 4 inches: cobbly loam
H2 - 4 to 30 inches: gravelly clay loam
H3 - 30 to 65 inches: channery clay loam

Properties and qualities

Slope: 0 to 8 percent
Percent of area covered with surface fragments: 9.0 percent
Depth to restrictive feature: 20 to 36 inches to fragipan
Natural drainage class: Moderately well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: About 18 to 36 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 3.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 7s
Hydrologic Soil Group: C
Hydric soil rating: No

Minor Components

Andover

Percent of map unit: 5 percent
Landform: Depressions
Landform position (three-dimensional): Mountainbase
Down-slope shape: Concave
Across-slope shape: Concave
Hydric soil rating: Yes

Philo

Percent of map unit: 3 percent
Hydric soil rating: No

Berks

Percent of map unit: 2 percent
Hydric soil rating: No

CbB—Clarksburg silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 16bq
Elevation: 200 to 1,500 feet
Mean annual precipitation: 32 to 48 inches
Mean annual air temperature: 48 to 57 degrees F
Frost-free period: 120 to 200 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Clarksburg and similar soils: 90 percent
Minor components: 5 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Clarksburg

Setting

Landform: Valley flats
Landform position (two-dimensional): Toeslope, footslope
Landform position (three-dimensional): Base slope
Down-slope shape: Linear, concave
Across-slope shape: Concave, linear
Parent material: Residuum weathered from limestone

Typical profile

Ap - 0 to 8 inches: silt loam
Bt - 8 to 27 inches: silt loam
Btx - 27 to 51 inches: silt loam
C - 51 to 84 inches: silt loam

Properties and qualities

Slope: 3 to 8 percent

Custom Soil Resource Report

Depth to restrictive feature: 20 to 36 inches to fragipan; 60 to 99 inches to
Natural drainage class: Moderately well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to
moderately high (0.06 to 0.60 in/hr)
Depth to water table: About 18 to 36 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Low (about 4.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 2e
Hydrologic Soil Group: C
Hydric soil rating: No

Minor Components

Thorndale

Percent of map unit: 5 percent
Landform: Depressions
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Base slope
Down-slope shape: Concave
Across-slope shape: Linear, concave
Hydric soil rating: Yes

EmB—Edom-Weikert complex, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 16c0
Elevation: 480 to 1,600 feet
Mean annual precipitation: 36 to 50 inches
Mean annual air temperature: 46 to 57 degrees F
Frost-free period: 120 to 210 days
Farmland classification: Farmland of statewide importance

Map Unit Composition

Edom and similar soils: 60 percent
Weikert and similar soils: 20 percent
Minor components: 5 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Edom

Setting

Landform: Hills
Landform position (two-dimensional): Summit
Landform position (three-dimensional): Side slope
Down-slope shape: Convex

Custom Soil Resource Report

Across-slope shape: Convex

Parent material: Clayey residuum weathered from limestone and shale

Typical profile

A - 0 to 8 inches: silty clay loam

Bt - 8 to 38 inches: silty clay

C - 38 to 46 inches: very channery silty clay loam

R - 46 to 50 inches: bedrock

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 40 to 72 inches to lithic bedrock

Natural drainage class: Well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.20 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum in profile: 10 percent

Salinity, maximum in profile: Nonsaline (0.0 to 0.2 mmhos/cm)

Available water storage in profile: Low (about 5.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: B

Hydric soil rating: No

Description of Weikert

Setting

Landform: Hillslopes

Landform position (two-dimensional): Shoulder

Landform position (three-dimensional): Side slope

Down-slope shape: Convex, linear

Across-slope shape: Convex, linear

Parent material: Residuum weathered from siltstone

Typical profile

A - 0 to 6 inches: channery silt loam

B - 6 to 15 inches: very channery silt loam

R - 15 to 19 inches: bedrock

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Natural drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Salinity, maximum in profile: Nonsaline (0.0 to 0.2 mmhos/cm)

Available water storage in profile: Very low (about 1.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Custom Soil Resource Report

Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: D
Hydric soil rating: No

Minor Components

Clarksburg

Percent of map unit: 5 percent
Hydric soil rating: No

MuB—Murrill gravelly silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 16dm
Elevation: 600 to 1,800 feet
Mean annual precipitation: 35 to 46 inches
Mean annual air temperature: 46 to 59 degrees F
Frost-free period: 120 to 180 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Murrill and similar soils: 90 percent
Minor components: 5 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Murrill

Setting

Landform: Valley sides
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Base slope
Down-slope shape: Concave
Across-slope shape: Concave
Parent material: Colluvium derived from sandstone over residuum weathered from limestone

Typical profile

H1 - 0 to 11 inches: channery silt loam
H2 - 11 to 60 inches: channery silty clay loam
H3 - 60 to 64 inches: gravelly sandy clay loam

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 72 to 99 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.20 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None

Custom Soil Resource Report

Frequency of ponding: None

Available water storage in profile: Moderate (about 7.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2e

Hydrologic Soil Group: B

Hydric soil rating: No

Minor Components

Buchanan

Percent of map unit: 5 percent

Hydric soil rating: No

MuD—Murrill gravelly silt loam, 15 to 25 percent slopes

Map Unit Setting

National map unit symbol: 16dp

Elevation: 400 to 3,800 feet

Mean annual precipitation: 34 to 60 inches

Mean annual air temperature: 48 to 57 degrees F

Frost-free period: 120 to 180 days

Farmland classification: Not prime farmland

Map Unit Composition

Murrill and similar soils: 90 percent

Minor components: 5 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Murrill

Setting

Landform: Valley sides

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave

Across-slope shape: Concave

Parent material: Colluvium derived from sandstone over residuum weathered from limestone

Typical profile

H1 - 0 to 11 inches: channery silt loam

H2 - 11 to 60 inches: channery silty clay loam

H3 - 60 to 64 inches: gravelly sandy clay loam

Properties and qualities

Slope: 15 to 25 percent

Depth to restrictive feature: 72 to 99 inches to lithic bedrock

Natural drainage class: Well drained

Runoff class: High

Custom Soil Resource Report

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.20 to 2.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Moderate (about 7.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: B

Hydric soil rating: No

Minor Components

Laidig

Percent of map unit: 5 percent

Hydric soil rating: No

MxB—Murrill extremely stony silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 16dq

Elevation: 600 to 2,400 feet

Mean annual precipitation: 35 to 46 inches

Mean annual air temperature: 46 to 57 degrees F

Frost-free period: 120 to 180 days

Farmland classification: Not prime farmland

Map Unit Composition

Murrill and similar soils: 90 percent

Minor components: 5 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Murrill

Setting

Landform: Valley sides

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave

Across-slope shape: Concave

Parent material: Colluvium derived from sandstone over residuum weathered from limestone

Typical profile

H1 - 0 to 11 inches: gravelly silt loam

H2 - 11 to 60 inches: channery silty clay loam

H3 - 60 to 64 inches: gravelly sandy clay loam

Custom Soil Resource Report

Properties and qualities

Slope: 3 to 8 percent
Percent of area covered with surface fragments: 9.0 percent
Depth to restrictive feature: 72 to 99 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.20 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Moderate (about 7.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 7s
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Buchanan

Percent of map unit: 5 percent
Hydric soil rating: No

OuC—Opequon silty clay loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 2sg9v
Elevation: 300 to 3,000 feet
Mean annual precipitation: 37 to 50 inches
Mean annual air temperature: 46 to 56 degrees F
Frost-free period: 155 to 192 days
Farmland classification: Farmland of statewide importance

Map Unit Composition

Opequon and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Opequon

Setting

Landform: Hills
Landform position (two-dimensional): Shoulder, summit, backslope
Landform position (three-dimensional): Nose slope, side slope, crest
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Clayey residuum weathered from limestone and dolomite

Custom Soil Resource Report

Typical profile

Ap - 0 to 5 inches: silty clay loam
Bt1 - 5 to 13 inches: silty clay
Bt2 - 13 to 16 inches: silty clay
R - 16 to 26 inches: bedrock

Properties and qualities

Slope: 8 to 15 percent
Depth to restrictive feature: 12 to 20 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.20 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline (0.0 to 0.2 mmhos/cm)
Available water storage in profile: Very low (about 1.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4s
Hydrologic Soil Group: D
Hydric soil rating: No

Minor Components

Hagerstown

Percent of map unit: 5 percent
Landform: Ridges
Landform position (two-dimensional): Backslope, footslope
Landform position (three-dimensional): Side slope, base slope
Down-slope shape: Linear
Across-slope shape: Linear
Hydric soil rating: No

Rock outcrop

Percent of map unit: 5 percent
Landform: Hills
Landform position (two-dimensional): Shoulder, summit, backslope
Landform position (three-dimensional): Nose slope, side slope, crest
Down-slope shape: Convex
Across-slope shape: Convex
Hydric soil rating: No

OuD—Opequon silty clay loam, 15 to 25 percent slopes

Map Unit Setting

National map unit symbol: 2sg9t
Elevation: 300 to 3,000 feet

Custom Soil Resource Report

Mean annual precipitation: 39 to 50 inches
Mean annual air temperature: 47 to 56 degrees F
Frost-free period: 155 to 192 days
Farmland classification: Not prime farmland

Map Unit Composition

Opequon and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Opequon

Setting

Landform: Hills
Landform position (two-dimensional): Shoulder, summit, backslope
Landform position (three-dimensional): Nose slope, side slope
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Clayey residuum weathered from limestone and dolomite

Typical profile

Ap - 0 to 5 inches: silty clay loam
Bt1 - 5 to 13 inches: silty clay
Bt2 - 13 to 16 inches: silty clay
R - 16 to 26 inches: bedrock

Properties and qualities

Slope: 15 to 25 percent
Depth to restrictive feature: 12 to 20 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.20 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline (0.0 to 0.2 mmhos/cm)
Available water storage in profile: Very low (about 1.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 6s
Hydrologic Soil Group: D
Hydric soil rating: No

Minor Components

Rock outcrop

Percent of map unit: 5 percent
Landform: Hills
Landform position (two-dimensional): Shoulder, summit, backslope
Landform position (three-dimensional): Nose slope, side slope, crest
Down-slope shape: Convex
Across-slope shape: Convex
Hydric soil rating: No

Hagerstown

Percent of map unit: 5 percent
Landform: Ridges
Landform position (two-dimensional): Backslope, footslope
Landform position (three-dimensional): Side slope, base slope
Down-slope shape: Linear
Across-slope shape: Linear
Hydric soil rating: No

UD—Udifluents-Dystrochrepts complex

Map Unit Setting

National map unit symbol: 16f0
Elevation: 200 to 1,300 feet
Mean annual precipitation: 30 to 50 inches
Mean annual air temperature: 45 to 55 degrees F
Frost-free period: 120 to 214 days
Farmland classification: Not prime farmland

Map Unit Composition

Udifluents and similar soils: 50 percent
Dystrochrepts and similar soils: 30 percent
Minor components: 4 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Udifluents

Setting

Landform: Flood plains
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Base slope
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Alluvium

Typical profile

H1 - 0 to 6 inches: channery silt loam
H2 - 6 to 42 inches: gravelly loam
H3 - 42 to 60 inches: silt loam

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Moderately well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.60 to 2.00 in/hr)
Depth to water table: About 36 inches
Frequency of flooding: Frequent
Frequency of ponding: None

Custom Soil Resource Report

Available water storage in profile: Moderate (about 6.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7s

Hydrologic Soil Group: C

Hydric soil rating: No

Description of Dystrochrepts

Setting

Landform: Mountain slopes

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Mountainflank

Down-slope shape: Linear, convex

Across-slope shape: Linear, convex

Parent material: Residuum weathered from sandstone and shale

Typical profile

H1 - 0 to 6 inches: loam, flagstones

H2 - 6 to 42 inches: gravelly loam

H3 - 42 to 60 inches: silt loam

Properties and qualities

Slope: 0 to 3 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Moderately well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.60 to 2.00 in/hr)

Depth to water table: About 36 inches

Frequency of flooding: Frequent

Frequency of ponding: None

Available water storage in profile: Low (about 6.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7s

Hydrologic Soil Group: C

Hydric soil rating: No

Minor Components

Brinkerton

Percent of map unit: 2 percent

Landform: Hills

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave

Across-slope shape: Concave

Hydric soil rating: Yes

Holly

Percent of map unit: 2 percent

Landform: Depressions on flood plains, backswamps

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave

Custom Soil Resource Report

Across-slope shape: Linear
Hydric soil rating: Yes

Huntingdon County, Pennsylvania

Ba—Barbour soils

Map Unit Setting

National map unit symbol: 15yf
Elevation: 400 to 800 feet
Mean annual precipitation: 32 to 50 inches
Mean annual air temperature: 45 to 55 degrees F
Frost-free period: 110 to 180 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Barbour and similar soils: 90 percent
Minor components: 7 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Barbour

Setting

Landform: Flood plains
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Reddish alluvium derived from sedimentary rock

Typical profile

Ap - 0 to 6 inches: fine sandy loam
B - 6 to 24 inches: sandy loam
2C - 24 to 60 inches: very gravelly loamy sand

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Natural drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.60 to 2.00 in/hr)
Depth to water table: About 36 to 72 inches
Frequency of flooding: Occasional
Frequency of ponding: None
Available water storage in profile: Low (about 5.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 1
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Basher

Percent of map unit: 5 percent
Hydric soil rating: No

Atkins

Percent of map unit: 2 percent

Custom Soil Resource Report

Landform: Flood plains
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Base slope
Down-slope shape: Concave
Across-slope shape: Concave
Hydric soil rating: Yes

HeD—Hagerstown-Rock outcrop complex, 5 to 25 percent slopes

Map Unit Setting

National map unit symbol: 15zt
Elevation: 460 to 1,500 feet
Mean annual precipitation: 30 to 46 inches
Mean annual air temperature: 45 to 59 degrees F
Frost-free period: 139 to 210 days
Farmland classification: Not prime farmland

Map Unit Composition

Hagerstown and similar soils: 75 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hagerstown

Setting

Landform: Ridges, valley floors
Landform position (two-dimensional): Backslope, shoulder
Landform position (three-dimensional): Side slope
Down-slope shape: Linear, convex
Across-slope shape: Linear, convex
Parent material: Residuum weathered from limestone

Typical profile

A - 0 to 8 inches: silt loam
Bt - 8 to 70 inches: clay

Properties and qualities

Slope: 5 to 25 percent
Depth to restrictive feature: 48 to 84 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: High (about 10.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4e

Custom Soil Resource Report

Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Hublersburg

Percent of map unit: 5 percent
Landform: Ridges on valleys
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear
Hydric soil rating: No

Edom

Percent of map unit: 5 percent
Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Hydric soil rating: No

OpD3—Opequon clay loam, 15 to 25 percent slopes, eroded

Map Unit Setting

National map unit symbol: 160y
Elevation: 400 to 3,000 feet
Mean annual precipitation: 30 to 45 inches
Mean annual air temperature: 45 to 57 degrees F
Frost-free period: 120 to 200 days
Farmland classification: Not prime farmland

Map Unit Composition

Opequon and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Opequon

Setting

Landform: Hills
Landform position (two-dimensional): Backslope, shoulder, summit
Landform position (three-dimensional): Side slope, crest
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Residuum weathered from limestone

Typical profile

A - 0 to 6 inches: clay loam
Bt - 6 to 18 inches: clay

Custom Soil Resource Report

R - 18 to 22 inches: bedrock

Properties and qualities

Slope: 15 to 25 percent

Depth to restrictive feature: 12 to 20 inches to lithic bedrock

Natural drainage class: Well drained

Runoff class: High

*Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.20 to 1.98 in/hr)*

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Very low (about 2.8 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6e

Hydrologic Soil Group: D

Hydric soil rating: No

Minor Components

Hagerstown

Percent of map unit: 5 percent

Hydric soil rating: No

Murrill

Percent of map unit: 5 percent

Hydric soil rating: No

Soil Information for All Uses

Soil Reports

The Soil Reports section includes various formatted tabular and narrative reports (tables) containing data for each selected soil map unit and each component of each unit. No aggregation of data has occurred as is done in reports in the Soil Properties and Qualities and Suitabilities and Limitations sections.

The reports contain soil interpretive information as well as basic soil properties and qualities. A description of each report (table) is included.

Land Classifications

This folder contains a collection of tabular reports that present a variety of soil groupings. The reports (tables) include all selected map units and components for each map unit. Land classifications are specified land use and management groupings that are assigned to soil areas because combinations of soil have similar behavior for specified practices. Most are based on soil properties and other factors that directly influence the specific use of the soil. Example classifications include ecological site classification, farmland classification, irrigated and nonirrigated land capability classification, and hydric rating.

Hydric Soils

This table lists the map unit components that are rated as hydric soils in the survey area. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt and others, 2002).

The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for all of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part

Custom Soil Resource Report

(Federal Register, 1994). These soils, under natural conditions, are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 2002). These criteria are used to identify map unit components that normally are associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (Soil Survey Staff, 1999) and "Keys to Soil Taxonomy" (Soil Survey Staff, 2006) and in the "Soil Survey Manual" (Soil Survey Division Staff, 1993).

If soils are wet enough for a long enough period of time to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils are specified in "Field Indicators of Hydric Soils in the United States" (Hurt and Vasilas, 2006).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

Map units that are dominantly made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units dominantly made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

The criteria for hydric soils are represented by codes in the table (for example, 2). Definitions for the codes are as follows:

1. All Histels except for Folistels, and Histosols except for Folists.
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Historthels great group, Histoturbels great group, Pachic subgroups, or Cumulic subgroups that:
 - A. Based on the range of characteristics for the soil series, will at least in part meet one or more Field Indicators of Hydric Soils in the United States, or
 - B. Show evidence that the soil meets the definition of a hydric soil;
3. Soils that are frequently ponded for long or very long duration during the growing season.
 - A. Based on the range of characteristics for the soil series, will at least in part meet one or more Field Indicators of Hydric Soils in the United States, or
 - B. Show evidence that the soil meets the definition of a hydric soil;
4. Map unit components that are frequently flooded for long duration or very long duration during the growing season that:
 - A. Based on the range of characteristics for the soil series, will at least in part meet one or more Field Indicators of Hydric Soils in the United States, or
 - B. Show evidence that the soil meets the definition of a hydric soil;

Custom Soil Resource Report

Hydric Condition: Food Security Act information regarding the ability to grow a commodity crop without removing woody vegetation or manipulating hydrology.

References:

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. September 18, 2002. Hydric soils of the United States.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18.

Soil Survey Staff. 2006. Keys to soil taxonomy. 10th edition. U.S. Department of Agriculture, Natural Resources Conservation Service.

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service. U.S. Department of Agriculture Handbook 436.

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

Report—Hydric Soils

Hydric Soils—Blair County, Pennsylvania				
Map symbol and map unit name	Component	Percent of map unit	Landform	Hydric criteria
BrB—Brinkerton silt loam, 3 to 8 percent slopes				
	Brinkerton	75	Depressions	2
	Atkins	3	Flood plains	2
BuB—Buchanan gravelly silt loam, 3 to 8 percent slopes				
	Andover	6	Depressions	2
BxB—Buchanan extremely stony silt loam, 3 to 8 percent slopes				
	Andover	5	Depressions	2
CbB—Clarksburg silt loam, 3 to 8 percent slopes				
	Thorndale	5	Depressions	2
UD—Udifluvents-Dystrochrepts complex				
	Brinkerton	2	Hills	2
	Holly	2	Depressions on flood plains, backswamps	2

Custom Soil Resource Report

Hydric Soils—Huntingdon County, Pennsylvania				
Map symbol and map unit name	Component	Percent of map unit	Landform	Hydric criteria
Ba—Barbour soils				
	Atkins	2	Flood plains	2

Vegetative Productivity

This folder contains a collection of tabular reports that present vegetative productivity data. The reports (tables) include all selected map units and components for each map unit. Vegetative productivity includes estimates of potential vegetative production for a variety of land uses, including cropland, forestland, hayland, pastureland, horticulture and rangeland. In the underlying database, some states maintain crop yield data by individual map unit component. Other states maintain the data at the map unit level. Attributes are included for both, although only one or the other is likely to contain data for any given geographic area. For other land uses, productivity data is shown only at the map unit component level. Examples include potential crop yields under irrigated and nonirrigated conditions, forest productivity, forest site index, and total rangeland production under of normal, favorable and unfavorable conditions.

Nonirrigated Yields by Map Unit

The average yields per acre that can be expected of the principal crops under a high level of management are shown in this table. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

If yields of irrigated crops are given, it is assumed that the irrigation system is adapted to the soils and to the crops grown, that good-quality irrigation water is uniformly applied as needed, and that tillage is kept to a minimum.

Pasture yields are expressed in terms of animal unit months. An animal unit month (AUM) is the amount of forage required by one mature cow of approximately 1,000 pounds weight, with or without a calf, for 1 month.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is

Custom Soil Resource Report

developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in the table are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local office of the Natural Resources Conservation Service or of the Cooperative Extension Service can provide information about the management and productivity of the soils for those crops.

The land capability classification of map units in the survey area is shown in this table. This classification shows, in a general way, the suitability of soils for most kinds of field crops (United States Department of Agriculture, Soil Conservation Service, 1961). Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes.

In the capability system, soils are generally grouped at three levels: capability class, subclass, and unit.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

- Class 1 soils have slight limitations that restrict their use.
- Class 2 soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.
- Class 3 soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.
- Class 4 soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.
- Class 5 soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.
- Class 6 soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.
- Class 7 soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.
- Class 8 soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, *e*, *w*, *s*, or *c*, to the class numeral, for example, 2e. The letter *e* shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; *w* shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); *s* shows that the soil is limited mainly because it is shallow, droughty, or stony; and

Custom Soil Resource Report

c, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by *w*, *s*, or *c* because the soils in class 5 are subject to little or no erosion.

Capability units are soil groups within a subclass. The soils in a capability unit are enough alike to be suited to the same crops and pasture plants, to require similar management, and to have similar productivity. Capability units are generally designated by adding an Arabic numeral to the subclass symbol, for example, 2e-4 and 3e-6. These units are not given in all soil surveys.

Reference:

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210.

Report—Nonirrigated Yields by Map Unit

Custom Soil Resource Report

Nonirrigated Yields by Map Unit—Blair County, Pennsylvania						
Map symbol and soil name	Land capability	Alfalfa hay	Corn	Grass-legume hay	Oats	Soybeans
		<i>Tons</i>	<i>Bu</i>	<i>Tons</i>	<i>Bu</i>	<i>Bu</i>
BrB—Brinkerton silt loam, 3 to 8 percent slopes		—	90	2.50	—	25
Brinkerton	4w					
BuB—Buchanan gravelly silt loam, 3 to 8 percent slopes		3.50	100	3.00	—	35
Buchanan	2w					
BxB—Buchanan extremely stony silt loam, 3 to 8 percent slopes		—	—	—	—	—
Buchanan	7s					
CbB—Clarksburg silt loam, 3 to 8 percent slopes		3.50	100	3.00	—	30
Clarksburg	2e					
EmB—Edom-Weikert complex, 3 to 8 percent slopes		4.00	100	3.00	70	—
Edom Weikert	3e 3e					
MuB—Murrill gravelly silt loam, 3 to 8 percent slopes		4.50	120	3.50	75	—
Murrill	2e					
MuD—Murrill gravelly silt loam, 15 to 25 percent slopes		4.00	95	3.00	60	—
Murrill	4e					
MxB—Murrill extremely stony silt loam, 3 to 8 percent slopes		—	—	—	—	—
Murrill	7s					
OuC—Opequon silty clay loam, 8 to 15 percent slopes		4.00	100	2.50	60	30
Opequon	4s					
OuD—Opequon silty clay loam, 15 to 25 percent slopes		4.00	100	2.50	60	30
Opequon	6s					
UD—Udifluvents-Dystrochrepts complex		—	—	—	—	—
Udifluvents Dystrochrepts	7s 7s					

Custom Soil Resource Report

Nonirrigated Yields by Map Unit—Huntingdon County, Pennsylvania						
Map symbol and soil name	Land capability	Alfalfa hay	Corn	Grass-legume hay	Oats	Soybeans
		<i>Tons</i>	<i>Bu</i>	<i>Tons</i>	<i>Bu</i>	<i>Bu</i>
Ba—Barbour soils		4.50	120	—	80	—
Barbour	1					
HeD—Hagerstown-Rock outcrop complex, 5 to 25 percent slopes		4.00	110	3.00	65	—
Hagerstown	4e					
OpD3—Opequon clay loam, 15 to 25 percent slopes, eroded		—	—	2.00	—	—
Opequon	6e					