



United States
Department of
Agriculture

NRCS

Natural
Resources
Conservation
Service

A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for Centre County, Pennsylvania



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<https://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	8
Soil Map.....	9
Legend.....	10
Map Unit Legend.....	11
Map Unit Descriptions.....	11
Centre County, Pennsylvania.....	14
HaB—Hagerstown silt loam, 3 to 8 percent slopes.....	14
HaC—Hagerstown silt loam, 8 to 15 percent slopes.....	15
HcD—Hagerstown silty clay loam, 15 to 25 percent slopes.....	17
HuB—Hublersburg silt loam, 3 to 8 percent slopes.....	18
HuC—Hublersburg silt loam, 8 to 15 percent slopes.....	19
Lx—Lindside soils.....	20
MrB—Morrison sandy loam, 2 to 8 percent slopes.....	22
MrD—Morrison sandy loam, 15 to 25 percent slopes.....	23
MuC—Murrill channery silt loam, 8 to 15 percent slopes.....	24
No—Nolin silt loam, local alluvium, 0 to 5 percent slopes.....	25
OhB—Opequon-Hagerstown complex, 3 to 8 percent slopes.....	27
OhC—Opequon-Hagerstown complex, 8 to 15 percent slopes.....	29
OhD—Opequon-Hagerstown complex, 15 to 25 percent slopes.....	31
Soil Information for All Uses	33
Soil Reports.....	33
Land Classifications.....	33
Land Capability Classification (Land Classification).....	33
Vegetative Productivity.....	35
Nonirrigated Yields by Map Unit.....	36
References	40

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

Custom Soil Resource Report

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

Custom Soil Resource Report

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.


Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map




Soil Map may not be valid at this scale.



MAP LEGEND

Area of Interest (AOI)

 Area of Interest (AOI)




















Soils

 Soil Map Unit Polygons

 Soil Map Unit Lines


 Soil Map Unit Points

Special Point Features






-  Blowout
-  Borrow Pit
-  Clay Spot
-  Closed Depression
-  Gravel Pit
-  Gravelly Spot
-  Landfill
-  Lava Flow
-  Marsh or swamp
-  Mine or Quarry
-  Miscellaneous Water
-  Perennial Water
-  Rock Outcrop
-  Saline Spot
-  Sandy Spot
-  Severely Eroded Spot
-  Sinkhole
-  Slide or Slip
-  Sodic Spot

-  Spoil Area
-  Stony Spot
-  Very Stony Spot
-  Wet Spot
-  Other
-  Special Line Features


Water Features

 Streams and Canals

Transportation

-  Rails
-  Interstate Highways
-  US Routes
-  Major Roads
-  Local Roads

Background

 Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:20,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Centre County, Pennsylvania
 Survey Area Data: Version 13, Sep 19, 2016

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Oct 6, 2011—Oct 17, 2011

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Centre County, Pennsylvania (PA027)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
HaB	Hagerstown silt loam, 3 to 8 percent slopes	22.0	29.9%
HaC	Hagerstown silt loam, 8 to 15 percent slopes	1.9	2.6%
HcD	Hagerstown silty clay loam, 15 to 25 percent slopes	1.1	1.4%
HuB	Hublersburg silt loam, 3 to 8 percent slopes	11.0	15.0%
HuC	Hublersburg silt loam, 8 to 15 percent slopes	2.9	3.9%
Lx	Lindside soils	3.1	4.2%
MrB	Morrison sandy loam, 2 to 8 percent slopes	18.9	25.8%
MrD	Morrison sandy loam, 15 to 25 percent slopes	1.6	2.1%
MuC	Murrill channery silt loam, 8 to 15 percent slopes	1.0	1.4%
No	Nolin silt loam, local alluvium, 0 to 5 percent slopes	2.7	3.6%
OhB	Opequon-Hagerstown complex, 3 to 8 percent slopes	2.4	3.3%
OhC	Opequon-Hagerstown complex, 8 to 15 percent slopes	0.5	0.7%
OhD	Opequon-Hagerstown complex, 15 to 25 percent slopes	4.5	6.1%
Totals for Area of Interest		73.5	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made

Custom Soil Resource Report

up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

Custom Soil Resource Report

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Centre County, Pennsylvania

HaB—Hagerstown silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2rc98
Elevation: 600 to 1,750 feet
Mean annual precipitation: 37 to 45 inches
Mean annual air temperature: 45 to 55 degrees F
Frost-free period: 155 to 190 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Hagerstown and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hagerstown

Setting

Landform: Hills
Landform position (two-dimensional): Backslope, footslope, summit
Landform position (three-dimensional): Side slope, base slope, interfluve
Down-slope shape: Linear, concave
Across-slope shape: Linear, concave
Parent material: Clayey residuum weathered from limestone

Typical profile

Ap - 0 to 10 inches: silt loam
Bt1 - 10 to 21 inches: silty clay loam
Bt2 - 21 to 56 inches: silty clay
C - 56 to 73 inches: silty clay loam
R - 73 to 83 inches: bedrock

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 43 to 98 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Moderate (about 8.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 2e
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Opequon

Percent of map unit: 5 percent

Custom Soil Resource Report

Landform: Ridges

Landform position (two-dimensional): Shoulder, summit

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Linear, convex

Across-slope shape: Convex, linear

Hydric soil rating: No

Carbo

Percent of map unit: 5 percent

Landform: Hills

Landform position (two-dimensional): Summit, backslope, shoulder

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Linear, convex

Across-slope shape: Linear, convex

Hydric soil rating: No

Funkstown

Percent of map unit: 3 percent

Landform: Valley floors

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave

Across-slope shape: Concave, linear

Hydric soil rating: No

Timberville

Percent of map unit: 2 percent

Landform: Hills

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Head slope, base slope

Down-slope shape: Concave, linear

Across-slope shape: Convex, concave, linear

Hydric soil rating: No

HaC—Hagerstown silt loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 2tb03

Elevation: 600 to 1,750 feet

Mean annual precipitation: 32 to 45 inches

Mean annual air temperature: 41 to 65 degrees F

Frost-free period: 155 to 181 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Hagerstown and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hagerstown

Setting

Landform: Hillslopes
Landform position (two-dimensional): Backslope, shoulder
Landform position (three-dimensional): Side slope
Down-slope shape: Linear, convex, concave
Across-slope shape: Linear, convex
Parent material: Clayey residuum weathered from limestone and dolomite

Typical profile

Ap - 0 to 8 inches: silt loam
Bt1 - 8 to 19 inches: silty clay loam
Bt2 - 19 to 54 inches: silty clay
C - 54 to 71 inches: silty clay loam
R - 71 to 81 inches: bedrock

Properties and qualities

Slope: 8 to 15 percent
Depth to restrictive feature: 43 to 98 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Salinity, maximum in profile: Nonsaline (0.0 to 1.9 mmhos/cm)
Available water storage in profile: Moderate (about 7.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Carbo

Percent of map unit: 8 percent
Landform: Hills
Landform position (two-dimensional): Summit, backslope, shoulder
Landform position (three-dimensional): Crest, side slope
Down-slope shape: Linear, convex
Across-slope shape: Linear, convex
Hydric soil rating: No

Opequon

Percent of map unit: 5 percent
Landform: Ridges
Landform position (two-dimensional): Shoulder, summit
Landform position (three-dimensional): Side slope, crest
Down-slope shape: Linear, convex
Across-slope shape: Convex, linear
Hydric soil rating: No

Clarksburg

Percent of map unit: 2 percent
Landform: Hillslopes
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Base slope, head slope
Down-slope shape: Linear
Across-slope shape: Concave
Hydric soil rating: No

HcD—Hagerstown silty clay loam, 15 to 25 percent slopes

Map Unit Setting

National map unit symbol: 2tb0h
Elevation: 600 to 1,750 feet
Mean annual precipitation: 37 to 45 inches
Mean annual air temperature: 46 to 54 degrees F
Frost-free period: 155 to 181 days
Farmland classification: Not prime farmland

Map Unit Composition

Hagerstown and similar soils: 80 percent
Minor components: 20 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hagerstown

Setting

Landform: Hills
Landform position (two-dimensional): Backslope, footslope
Landform position (three-dimensional): Side slope, nose slope
Down-slope shape: Linear, concave
Across-slope shape: Linear, concave
Parent material: Clayey residuum weathered from limestone

Typical profile

Ap - 0 to 7 inches: silty clay loam
Bt1 - 7 to 19 inches: silty clay loam
Bt2 - 19 to 31 inches: silty clay
C - 31 to 59 inches: silty clay
R - 59 to 69 inches: bedrock

Properties and qualities

Slope: 15 to 25 percent
Depth to restrictive feature: 40 to 79 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None

Custom Soil Resource Report

Frequency of ponding: None

Salinity, maximum in profile: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water storage in profile: High (about 9.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: B

Other vegetative classification: Moist Loams (ML2)

Hydric soil rating: No

Minor Components

Carbo

Percent of map unit: 13 percent

Landform: Hills

Landform position (two-dimensional): Summit, backslope, shoulder

Landform position (three-dimensional): Crest, side slope

Down-slope shape: Linear, convex

Across-slope shape: Linear, convex

Hydric soil rating: No

Opequon

Percent of map unit: 7 percent

Landform: Hills

Landform position (two-dimensional): Shoulder, summit, backslope

Landform position (three-dimensional): Nose slope, side slope, crest

Down-slope shape: Convex

Across-slope shape: Convex

Hydric soil rating: No

HuB—Hublersburg silt loam, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: I249

Elevation: 310 to 3,000 feet

Mean annual precipitation: 30 to 50 inches

Mean annual air temperature: 45 to 57 degrees F

Frost-free period: 120 to 205 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Hublersburg and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hublersburg

Setting

Landform: Valleys

Custom Soil Resource Report

Landform position (two-dimensional): Summit
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear

Typical profile

Ap - 0 to 9 inches: silt loam
Bt - 9 to 70 inches: silty clay

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 60 to 99 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Moderate (about 8.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 2e
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Nolin

Percent of map unit: 5 percent
Hydric soil rating: No

Opequeon

Percent of map unit: 5 percent
Hydric soil rating: No

HuC—Hublersburg silt loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: I24b
Elevation: 310 to 3,000 feet
Mean annual precipitation: 30 to 50 inches
Mean annual air temperature: 45 to 57 degrees F
Frost-free period: 120 to 205 days
Farmland classification: Farmland of statewide importance

Map Unit Composition

Hublersburg and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hublersburg

Setting

Landform: Valleys
Landform position (two-dimensional): Summit
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear

Typical profile

Ap - 0 to 9 inches: silt loam
Bt - 9 to 70 inches: silty clay

Properties and qualities

Slope: 8 to 15 percent
Depth to restrictive feature: 60 to 99 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Moderate (about 8.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Opequeon

Percent of map unit: 7 percent
Hydric soil rating: No

Nolin

Percent of map unit: 3 percent
Hydric soil rating: No

Lx—Lindside soils

Map Unit Setting

National map unit symbol: I24w
Elevation: 300 to 1,500 feet
Mean annual precipitation: 32 to 55 inches
Mean annual air temperature: 46 to 57 degrees F
Frost-free period: 133 to 205 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Lindside and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Lindside

Setting

Landform: Flood plains

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Alluvium derived from limestone

Typical profile

H1 - 0 to 8 inches: silt loam

H2 - 8 to 44 inches: silt loam

H3 - 44 to 60 inches: loam

Properties and qualities

Slope: 0 to 3 percent

Depth to restrictive feature: 72 to 99 inches to

Natural drainage class: Moderately well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.20 to 2.00 in/hr)

Depth to water table: About 18 to 36 inches

Frequency of flooding: Occasional

Frequency of ponding: None

Available water storage in profile: High (about 11.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: C

Hydric soil rating: No

Minor Components

Chagrin

Percent of map unit: 5 percent

Landform: Flood plains

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Hydric soil rating: No

Melvin

Percent of map unit: 5 percent

Landform: Flood plains

Down-slope shape: Concave

Across-slope shape: Concave

Hydric soil rating: Yes

MrB—Morrison sandy loam, 2 to 8 percent slopes

Map Unit Setting

National map unit symbol: I258
Elevation: 600 to 1,800 feet
Mean annual precipitation: 35 to 50 inches
Mean annual air temperature: 46 to 55 degrees F
Frost-free period: 120 to 180 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Morrison and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Morrison

Setting

Landform: Ridges
Landform position (two-dimensional): Backslope, summit
Landform position (three-dimensional): Crest, side slope
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Residuum weathered from limestone and sandstone

Typical profile

H1 - 0 to 14 inches: sandy loam
H2 - 14 to 53 inches: sandy loam
H3 - 53 to 74 inches: channery sandy loam

Properties and qualities

Slope: 2 to 8 percent
Depth to restrictive feature: 72 to 99 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 6.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Moderate (about 6.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 2e
Hydrologic Soil Group: A
Hydric soil rating: No

Minor Components

Vanderlip

Percent of map unit: 5 percent
Landform: Ridges
Landform position (two-dimensional): Summit, shoulder
Landform position (three-dimensional): Mountaintop, side slope
Down-slope shape: Convex
Across-slope shape: Convex
Hydric soil rating: No

Murrill

Percent of map unit: 5 percent
Hydric soil rating: No

MrD—Morrison sandy loam, 15 to 25 percent slopes

Map Unit Setting

National map unit symbol: I25b
Elevation: 600 to 1,800 feet
Mean annual precipitation: 35 to 50 inches
Mean annual air temperature: 46 to 55 degrees F
Frost-free period: 120 to 180 days
Farmland classification: Not prime farmland

Map Unit Composition

Morrison and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Morrison

Setting

Landform: Ridges
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Crest, side slope
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Residuum weathered from limestone and sandstone

Typical profile

H1 - 0 to 14 inches: sandy loam
H2 - 14 to 53 inches: sandy loam
H3 - 53 to 74 inches: channery sandy loam

Properties and qualities

Slope: 15 to 25 percent
Depth to restrictive feature: 72 to 99 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium

Custom Soil Resource Report

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Moderate (about 6.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: A

Hydric soil rating: No

Minor Components

Vanderlip

Percent of map unit: 5 percent

Landform: Ridges

Landform position (two-dimensional): Summit, shoulder

Landform position (three-dimensional): Mountaintop, side slope

Down-slope shape: Convex

Across-slope shape: Convex

Hydric soil rating: No

Murrill

Percent of map unit: 5 percent

Hydric soil rating: No

MuC—Murrill channery silt loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: I25h

Elevation: 400 to 3,000 feet

Mean annual precipitation: 30 to 45 inches

Mean annual air temperature: 45 to 57 degrees F

Frost-free period: 120 to 200 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Murrill and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Murrill

Setting

Landform: Valley sides

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave

Across-slope shape: Concave

Custom Soil Resource Report

Parent material: Colluvium derived from sandstone over residuum weathered from limestone

Typical profile

H1 - 0 to 15 inches: channery silt loam
H2 - 15 to 60 inches: channery clay loam
H3 - 60 to 80 inches: clay

Properties and qualities

Slope: 8 to 15 percent
Depth to restrictive feature: 72 to 99 inches to
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.20 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Moderate (about 7.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Hagerstown

Percent of map unit: 10 percent
Hydric soil rating: No

Opequon

Percent of map unit: 5 percent
Hydric soil rating: No

No—Nolin silt loam, local alluvium, 0 to 5 percent slopes

Map Unit Setting

National map unit symbol: I25m
Elevation: 300 to 1,500 feet
Mean annual precipitation: 34 to 50 inches
Mean annual air temperature: 46 to 57 degrees F
Frost-free period: 120 to 214 days
Farmland classification: All areas are prime farmland

Map Unit Composition

Nolin and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Nolin

Setting

Landform: Valleys
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Alluvium derived from sedimentary rock over residuum weathered from limestone and shale

Typical profile

H1 - 0 to 8 inches: silt loam
H2 - 8 to 52 inches: silt loam
H3 - 52 to 65 inches: silty clay loam

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: 72 to 99 inches to
Natural drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: About 36 to 72 inches
Frequency of flooding: Rare
Frequency of ponding: None
Available water storage in profile: Very high (about 12.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 1
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Clarksburg

Percent of map unit: 5 percent
Hydric soil rating: No

Melvin

Percent of map unit: 5 percent
Landform: Flood plains
Down-slope shape: Concave
Across-slope shape: Concave
Hydric soil rating: Yes

Dunning

Percent of map unit: 5 percent
Landform: Flood plains, terraces
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Tread
Down-slope shape: Concave
Across-slope shape: Concave
Hydric soil rating: Yes

OhB—Opequon-Hagerstown complex, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: I25p

Elevation: 400 to 3,000 feet

Mean annual precipitation: 30 to 46 inches

Mean annual air temperature: 45 to 57 degrees F

Frost-free period: 140 to 210 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Opequon and similar soils: 50 percent

Hagerstown and similar soils: 30 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Opequon

Setting

Landform: Hills

Landform position (two-dimensional): Backslope, shoulder, summit

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Residuum weathered from limestone

Typical profile

H1 - 0 to 6 inches: silty clay loam

H2 - 6 to 16 inches: silty clay

H3 - 16 to 20 inches: bedrock

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 12 to 20 inches to lithic bedrock

Natural drainage class: Well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high
(0.06 to 2.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Very low (about 2.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: D

Hydric soil rating: No

Description of Hagerstown

Setting

Landform: Hills
Landform position (two-dimensional): Summit
Landform position (three-dimensional): Crest
Down-slope shape: Convex
Across-slope shape: Convex

Typical profile

A - 0 to 8 inches: silt loam
Bt - 8 to 45 inches: clay
C - 45 to 75 inches: clay loam

Properties and qualities

Slope: 3 to 8 percent
Depth to restrictive feature: 40 to 72 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: High (about 9.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 2e
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Edom

Percent of map unit: 5 percent
Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Hydric soil rating: No

Hagerstown

Percent of map unit: 5 percent
Hydric soil rating: No

OhC—Opequon-Hagerstown complex, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 125q

Elevation: 400 to 3,000 feet

Mean annual precipitation: 30 to 46 inches

Mean annual air temperature: 45 to 57 degrees F

Frost-free period: 140 to 210 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Opequon and similar soils: 55 percent

Hagerstown and similar soils: 25 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Opequon

Setting

Landform: Hills

Landform position (two-dimensional): Backslope, shoulder, summit

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Residuum weathered from limestone

Typical profile

H1 - 0 to 6 inches: silty clay loam

H2 - 6 to 16 inches: silty clay

H3 - 16 to 20 inches: bedrock

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: 12 to 20 inches to lithic bedrock

Natural drainage class: Well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high
(0.06 to 2.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Very low (about 2.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: D

Hydric soil rating: No

Description of Hagerstown

Setting

Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear

Typical profile

A - 0 to 8 inches: silt loam
Bt - 8 to 45 inches: clay
C - 45 to 75 inches: clay loam

Properties and qualities

Slope: 8 to 15 percent
Depth to restrictive feature: 40 to 72 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: High (about 9.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 3e
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Edom

Percent of map unit: 5 percent
Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Hydric soil rating: No

Hagerstown

Percent of map unit: 5 percent
Hydric soil rating: No

OhD—Opequon-Hagerstown complex, 15 to 25 percent slopes

Map Unit Setting

National map unit symbol: 125r
Elevation: 400 to 3,000 feet
Mean annual precipitation: 30 to 46 inches
Mean annual air temperature: 45 to 57 degrees F
Frost-free period: 140 to 210 days
Farmland classification: Not prime farmland

Map Unit Composition

Opequon and similar soils: 60 percent
Hagerstown and similar soils: 20 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Opequon

Setting

Landform: Hills
Landform position (two-dimensional): Backslope, shoulder, summit
Landform position (three-dimensional): Side slope, crest
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Residuum weathered from limestone

Typical profile

H1 - 0 to 6 inches: silty clay loam
H2 - 6 to 16 inches: silty clay
H3 - 16 to 20 inches: bedrock

Properties and qualities

Slope: 15 to 25 percent
Depth to restrictive feature: 12 to 20 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high
(0.06 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: Very low (about 2.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 6e
Hydrologic Soil Group: D
Hydric soil rating: No

Description of Hagerstown

Setting

Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear

Typical profile

A - 0 to 8 inches: silt loam
Bt - 8 to 45 inches: clay
C - 45 to 75 inches: clay loam

Properties and qualities

Slope: 15 to 25 percent
Depth to restrictive feature: 40 to 72 inches to lithic bedrock
Natural drainage class: Well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water storage in profile: High (about 9.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4e
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Edom

Percent of map unit: 5 percent
Landform: Hillslopes
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Convex
Hydric soil rating: No

Hagerstown

Percent of map unit: 5 percent
Hydric soil rating: No

Soil Information for All Uses

Soil Reports

The Soil Reports section includes various formatted tabular and narrative reports (tables) containing data for each selected soil map unit and each component of each unit. No aggregation of data has occurred as is done in reports in the Soil Properties and Qualities and Suitabilities and Limitations sections.

The reports contain soil interpretive information as well as basic soil properties and qualities. A description of each report (table) is included.

Land Classifications

This folder contains a collection of tabular reports that present a variety of soil groupings. The reports (tables) include all selected map units and components for each map unit. Land classifications are specified land use and management groupings that are assigned to soil areas because combinations of soil have similar behavior for specified practices. Most are based on soil properties and other factors that directly influence the specific use of the soil. Example classifications include ecological site classification, farmland classification, irrigated and nonirrigated land capability classification, and hydric rating.

Land Capability Classification (Land Classification)

The land capability classification of map units in the survey area is shown in this table. This classification shows, in a general way, the suitability of soils for most kinds of field crops (United States Department of Agriculture, Soil Conservation Service, 1961). Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes.

In the capability system, soils are generally grouped at three levels: capability class, subclass, and unit.

Custom Soil Resource Report

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

- Class 1 soils have slight limitations that restrict their use.
- Class 2 soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.
- Class 3 soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.
- Class 4 soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.
- Class 5 soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.
- Class 6 soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.
- Class 7 soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.
- Class 8 soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, *e*, *w*, *s*, or *c*, to the class numeral, for example, 2e. The letter *e* shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; *w* shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); *s* shows that the soil is limited mainly because it is shallow, droughty, or stony; and *c*, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by *w*, *s*, or *c* because the soils in class 5 are subject to little or no erosion.

Report—Land Capability Classification (Land Classification)

Land Capability Classification—Centre County, Pennsylvania				
Map unit symbol and name	Pct. of map unit	Component name	Land Capability Subclass	
			Nonirrigated	Irrigated
HaB—Hagerstown silt loam, 3 to 8 percent slopes				
	85	Hagerstown	2e	—
HaC—Hagerstown silt loam, 8 to 15 percent slopes				
	85	Hagerstown	3e	—

Custom Soil Resource Report

Land Capability Classification—Centre County, Pennsylvania				
Map unit symbol and name	Pct. of map unit	Component name	Land Capability Subclass	
			Nonirrigated	Irrigated
HcD—Hagerstown silty clay loam, 15 to 25 percent slopes				
	80	Hagerstown	4e	—
HuB—Hublersburg silt loam, 3 to 8 percent slopes				
	90	Hublersburg	2e	—
HuC—Hublersburg silt loam, 8 to 15 percent slopes				
	90	Hublersburg	3e	—
Lx—Lindside soils				
	90	Lindside	2w	—
MrB—Morrison sandy loam, 2 to 8 percent slopes				
	90	Morrison	2e	—
MrD—Morrison sandy loam, 15 to 25 percent slopes				
	90	Morrison	4e	—
MuC—Murrill channery silt loam, 8 to 15 percent slopes				
	85	Murrill	3e	—
No—Nolin silt loam, local alluvium, 0 to 5 percent slopes				
	85	Nolin	1	—
OhB—Opequon-Hagerstown complex, 3 to 8 percent slopes				
	50	Opequon	3e	—
	30	Hagerstown	2e	—
OhC—Opequon-Hagerstown complex, 8 to 15 percent slopes				
	55	Opequon	4e	—
	25	Hagerstown	3e	—
OhD—Opequon-Hagerstown complex, 15 to 25 percent slopes				
	60	Opequon	6e	—
	20	Hagerstown	4e	—

Vegetative Productivity

This folder contains a collection of tabular reports that present vegetative productivity data. The reports (tables) include all selected map units and

components for each map unit. Vegetative productivity includes estimates of potential vegetative production for a variety of land uses, including cropland, forestland, hayland, pastureland, horticulture and rangeland. In the underlying database, some states maintain crop yield data by individual map unit component. Other states maintain the data at the map unit level. Attributes are included for both, although only one or the other is likely to contain data for any given geographic area. For other land uses, productivity data is shown only at the map unit component level. Examples include potential crop yields under irrigated and nonirrigated conditions, forest productivity, forest site index, and total rangeland production under of normal, favorable and unfavorable conditions.

Nonirrigated Yields by Map Unit

The average yields per acre that can be expected of the principal crops under a high level of management are shown in this table. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

If yields of irrigated crops are given, it is assumed that the irrigation system is adapted to the soils and to the crops grown, that good-quality irrigation water is uniformly applied as needed, and that tillage is kept to a minimum.

Pasture yields are expressed in terms of animal unit months. An animal unit month (AUM) is the amount of forage required by one mature cow of approximately 1,000 pounds weight, with or without a calf, for 1 month.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in the table are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local office of the Natural Resources Conservation Service or of the Cooperative Extension Service can provide information about the management and productivity of the soils for those crops.

The land capability classification of map units in the survey area is shown in this table. This classification shows, in a general way, the suitability of soils for most kinds of field crops (United States Department of Agriculture, Soil Conservation Service, 1961). Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in

Custom Soil Resource Report

grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes.

In the capability system, soils are generally grouped at three levels: capability class, subclass, and unit.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

- Class 1 soils have slight limitations that restrict their use.
- Class 2 soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.
- Class 3 soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.
- Class 4 soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.
- Class 5 soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.
- Class 6 soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.
- Class 7 soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.
- Class 8 soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, *e*, *w*, *s*, or *c*, to the class numeral, for example, 2*e*. The letter *e* shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; *w* shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); *s* shows that the soil is limited mainly because it is shallow, droughty, or stony; and *c*, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by *w*, *s*, or *c* because the soils in class 5 are subject to little or no erosion.

Capability units are soil groups within a subclass. The soils in a capability unit are enough alike to be suited to the same crops and pasture plants, to require similar management, and to have similar productivity. Capability units are generally designated by adding an Arabic numeral to the subclass symbol, for example, 2*e*-4 and 3*e*-6. These units are not given in all soil surveys.

Reference:

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210.

Report—Nonirrigated Yields by Map Unit

Custom Soil Resource Report

Nonirrigated Yields by Map Unit—Centre County, Pennsylvania						
Map symbol and soil name	Land capability	Alfalfa hay	Corn	Corn silage	Grass-legume hay	Soybeans
		<i>Tons</i>	<i>Bu</i>	<i>Tons</i>	<i>Tons</i>	<i>Bu</i>
HaB—Hagerstown silt loam, 3 to 8 percent slopes		6.00	155	25.00	3.50	50
Hagerstown	2e					
HaC—Hagerstown silt loam, 8 to 15 percent slopes		5.60	144	23.00	3.50	47
Hagerstown	3e					
HcD—Hagerstown silty clay loam, 15 to 25 percent slopes		4.80	124	20.00	2.80	—
Hagerstown	4e					
HuB—Hublersburg silt loam, 3 to 8 percent slopes		5.00	130	26.00	3.50	—
Hublersburg	2e					
HuC—Hublersburg silt loam, 8 to 15 percent slopes		4.50	125	25.00	3.00	—
Hublersburg	3e					
Lx—Lindside soils		4.50	125	—	3.50	45
Lindside	2w					
MrB—Morrison sandy loam, 2 to 8 percent slopes		—	100	20.00	3.50	—
Morrison	2e					
MrD—Morrison sandy loam, 15 to 25 percent slopes		—	90	18.00	3.00	—
Morrison	4e					
MuC—Murrill channery silt loam, 8 to 15 percent slopes		4.00	110	22.00	3.00	—
Murrill	3e					
No—Nolin silt loam, local alluvium, 0 to 5 percent slopes		—	135	—	4.50	45
Nolin	1					
OhB—Opequon-Hagerstown complex, 3 to 8 percent slopes		—	75	—	2.50	—
Opequon Hagerstown	3e 2e					
OhC—Opequon-Hagerstown complex, 8 to 15 percent slopes		—	70	—	2.50	—
Opequon Hagerstown	4e 3e					
OhD—Opequon-Hagerstown complex, 15 to 25 percent slopes		—	—	—	2.00	—
Opequon Hagerstown	6e 4e					

References

- American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.
- American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.
- Federal Register. July 13, 1994. Changes in hydric soils of the United States.
- Federal Register. September 18, 2002. Hydric soils of the United States.
- Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.
- National Research Council. 1995. Wetlands: Characteristics and boundaries.
- Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_054262
- Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577
- Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580
- Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.
- United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.
- United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053374
- United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. <http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084>

Custom Soil Resource Report

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf