


Natural Resources Conservation

Service

A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

# Custom Soil Resource Report for Huntingdon County, Pennsylvania

**Ormsby Dairy Farm** 



# **Preface**

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2 053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.


# **Contents**

Preface	2
Soil Map	5
Soil Map	6
Legend	7
Map Unit Legend	
Map Unit Descriptions	8
Huntingdon County, Pennsylvania	11
BuB—Buchanan gravelly loam, 3 to 8 percent slopes	11
EgC—Edom-Weikert complex, 8 to 15 percent slopes	12
EgD—Edom-Weikert complex, 15 to 25 percent slopes	14
EgF—Edom-Weikert complex, 25 to 60 percent slopes	16
HeD—Hagerstown-Rock outcrop complex, 5 to 25 percent slopes	17
MoB—Monongahela silt loam, 2 to 10 percent slopes	19
MuB—Murrill gravelly loam, 3 to 8 percent slopes	20
MuC—Murrill gravelly loam, 8 to 15 percent slopes	21
Ne—Newark silt loam	23
Po—Philo and Basher silt loams, high bottom	24
Pu—Purdy silt loam	26
W—Water	27
Soil Information for All Uses	28
Soil Reports	28
Vegetative Productivity	28
Nonirrigated Yields by Map Unit	28
References	32

# Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

# Custom Soil Resource Report Soil Map



#### MAP LEGEND

#### Area of Interest (AOI)

Area of Interest (AOI)

#### Soils

Soil Map Unit Polygons

Soil Map Unit Lines



Soil Map Unit Points

#### **Special Point Features**

Blowout (o)

Borrow Pit

Clay Spot

**Closed Depression** 

Gravel Pit

**Gravelly Spot** 

Landfill

Lava Flow Marsh or swamp

Mine or Quarry

Miscellaneous Water

Perennial Water Rock Outcrop

Saline Spot

Sandy Spot

Severely Eroded Spot

Sinkhole

Sodic Spot

Slide or Slip



Spoil Area Stony Spot



Very Stony Spot



Wet Spot Other



Special Line Features

### **Water Features**

Streams and Canals

#### Transportation

---

Rails

Interstate Highways

**US Routes** 

Major Roads

00

Local Roads

### Background

Aerial Photography

#### MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:20.000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service Web Soil Survey URL:

Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Huntingdon County, Pennsylvania Survey Area Data: Version 14, Jun 5, 2020

Soil map units are labeled (as space allows) for map scales 1:50.000 or larger.

Date(s) aerial images were photographed: Oct 6, 2011—Oct 17, 2011

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

# Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI				
BuB	Buchanan gravelly loam, 3 to 8 percent slopes	10.2	6.9%				
EgC	Edom-Weikert complex, 8 to 15 percent slopes	20.7	14.0%				
EgD	Edom-Weikert complex, 15 to 25 percent slopes	11.1	7.5%				
EgF	Edom-Weikert complex, 25 to 60 percent slopes	20.3	13.7%				
HeD	Hagerstown-Rock outcrop complex, 5 to 25 percent slopes	0.2	0.1%				
МоВ	Monongahela silt loam, 2 to 10 percent slopes	46.3	31.4%				
MuB	Murrill gravelly loam, 3 to 8 percent slopes	18.2	12.3%				
MuC	Murrill gravelly loam, 8 to 15 percent slopes	11.7	7.9%				
Ne	Newark silt loam	1.9	1.3%				
Ро	Philo and Basher silt loams, high bottom	4.4	3.0%				
Pu	Purdy silt loam	2.6	1.7%				
W	Water	0.0	0.0%				
Totals for Area of Interest		147.7	100.0%				

# **Map Unit Descriptions**

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called

noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can

be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

### **Huntingdon County, Pennsylvania**

### BuB—Buchanan gravelly loam, 3 to 8 percent slopes

### **Map Unit Setting**

National map unit symbol: 15yw Elevation: 400 to 3,800 feet

Mean annual precipitation: 34 to 60 inches Mean annual air temperature: 46 to 59 degrees F

Frost-free period: 110 to 180 days

Farmland classification: All areas are prime farmland

### **Map Unit Composition**

Buchanan and similar soils: 75 percent

Minor components: 25 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Buchanan**

### Setting

Landform: Valley sides, mountain slopes

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Lower third of mountainflank

Down-slope shape: Concave, linear Across-slope shape: Concave, linear

Parent material: Colluvium derived from sandstone and shale

### Typical profile

Ap - 0 to 7 inches: gravelly loam

Bt - 7 to 21 inches: gravelly loam

Btx - 21 to 65 inches: cobbly clay loam

### Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: 20 to 36 inches to fragipan; 60 to 99 inches to lithic

bedrock

Drainage class: Moderately well drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to

moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 14 to 30 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Very low (about 2.9 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2e

Hydrologic Soil Group: D Hydric soil rating: No

### **Minor Components**

### **Andover**

Percent of map unit: 10 percent

Landform: Depressions

Landform position (two-dimensional): Footslope, toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave Hydric soil rating: Yes

Wharton

Percent of map unit: 10 percent

Landform: Hills

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Interfluve, side slope, head slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: No

Laidig

Percent of map unit: 5 percent

Landform: Mountains

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Lower third of mountainflank

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: No

### EgC—Edom-Weikert complex, 8 to 15 percent slopes

### **Map Unit Setting**

National map unit symbol: 15zf Elevation: 300 to 3.000 feet

Mean annual precipitation: 30 to 50 inches
Mean annual air temperature: 45 to 57 degrees F

Frost-free period: 120 to 210 days

Farmland classification: Farmland of statewide importance

### **Map Unit Composition**

Edom and similar soils: 60 percent Weikert and similar soils: 20 percent Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Edom**

Setting

Landform: Hills

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Convex

Parent material: Clayey residuum weathered from limestone and shale

### Typical profile

H1 - 0 to 8 inches: silty clay loam H2 - 8 to 38 inches: silty clay

H3 - 38 to 46 inches: very channery silty clay loam

H4 - 46 to 50 inches: bedrock

### Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: 40 to 72 inches to lithic bedrock

Drainage class: Well drained Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.20 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Calcium carbonate, maximum content: 10 percent Maximum salinity: Nonsaline (0.0 to 0.2 mmhos/cm) Available water capacity: Low (about 5.4 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: B Hydric soil rating: No

### **Description of Weikert**

### Setting

Landform: Hillslopes

Landform position (two-dimensional): Shoulder Landform position (three-dimensional): Side slope

Down-slope shape: Convex, linear Across-slope shape: Convex, linear

Parent material: Residuum weathered from siltstone

### **Typical profile**

H1 - 0 to 6 inches: channery silt loam
H2 - 6 to 15 inches: very channery silt loam

H3 - 15 to 19 inches: bedrock

### **Properties and qualities**

Slope: 8 to 15 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 0.2 mmhos/cm) Available water capacity: Very low (about 1.2 inches)

#### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: D Hydric soil rating: No

### **Minor Components**

### Opequon

Percent of map unit: 5 percent Hydric soil rating: No

### Clarksburg

Percent of map unit: 5 percent Hydric soil rating: No

### EgD—Edom-Weikert complex, 15 to 25 percent slopes

### **Map Unit Setting**

National map unit symbol: 15zg Elevation: 300 to 3,000 feet

Mean annual precipitation: 30 to 50 inches Mean annual air temperature: 45 to 57 degrees F

Frost-free period: 120 to 210 days

Farmland classification: Not prime farmland

### **Map Unit Composition**

Edom and similar soils: 60 percent Weikert and similar soils: 20 percent Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Edom**

### Setting

Landform: Hills

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Convex

Parent material: Clayey residuum weathered from limestone and shale

### **Typical profile**

H1 - 0 to 8 inches: silty clay loam H2 - 8 to 38 inches: silty clay

H3 - 38 to 46 inches: very channery silty clay loam

H4 - 46 to 50 inches: bedrock

### **Properties and qualities**

Slope: 15 to 25 percent

Depth to restrictive feature: 40 to 72 inches to lithic bedrock

Drainage class: Well drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.20 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Calcium carbonate, maximum content: 10 percent Maximum salinity: Nonsaline (0.0 to 0.2 mmhos/cm) Available water capacity: Low (about 5.4 inches)

#### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6e

Hydrologic Soil Group: B Hydric soil rating: No

### **Description of Weikert**

### Setting

Landform: Hillslopes

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex, linear Across-slope shape: Convex, linear

Parent material: Residuum weathered from siltstone

### Typical profile

H1 - 0 to 6 inches: channery silt loam H2 - 6 to 15 inches: very channery silt loam

H3 - 15 to 19 inches: bedrock

### **Properties and qualities**

Slope: 15 to 25 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Somewhat excessively drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 0.2 mmhos/cm) Available water capacity: Very low (about 1.2 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6e

Hydrologic Soil Group: D Hydric soil rating: No

### **Minor Components**

#### Klinesville

Percent of map unit: 5 percent Hydric soil rating: No

### Opequon

Percent of map unit: 5 percent

Hydric soil rating: No

### EgF—Edom-Weikert complex, 25 to 60 percent slopes

### **Map Unit Setting**

National map unit symbol: 15zh Elevation: 300 to 3,000 feet

Mean annual precipitation: 30 to 50 inches Mean annual air temperature: 45 to 57 degrees F

Frost-free period: 120 to 210 days

Farmland classification: Not prime farmland

### **Map Unit Composition**

Edom and similar soils: 50 percent Weikert and similar soils: 30 percent Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Edom**

### Setting

Landform: Hills

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Convex

Parent material: Residuum weathered from limestone and shale

### Typical profile

H1 - 0 to 8 inches: silty clay loam H2 - 8 to 38 inches: silty clay

H3 - 38 to 46 inches: very channery silty clay loam

H4 - 46 to 50 inches: bedrock

### **Properties and qualities**

Slope: 25 to 60 percent

Depth to restrictive feature: 40 to 72 inches to lithic bedrock

Drainage class: Well drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.20 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Low (about 5.4 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7e

Hydrologic Soil Group: B Hydric soil rating: No

### **Description of Weikert**

### Setting

Landform: Hills

Landform position (two-dimensional): Backslope, shoulder Landform position (three-dimensional): Side slope, crest

Down-slope shape: Convex Across-slope shape: Convex

Parent material: Residuum weathered from shale and siltstone

### **Typical profile**

H1 - 0 to 3 inches: channery silt loam H2 - 3 to 15 inches: very channery silt loam

H3 - 15 to 19 inches: bedrock

### Properties and qualities

Slope: 25 to 50 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Well drained Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Very low (about 1.1 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6e

Hydrologic Soil Group: D Hydric soil rating: No

### **Minor Components**

### Klinesville

Percent of map unit: 5 percent

Hydric soil rating: No

### Opequon

Percent of map unit: 5 percent

Hydric soil rating: No

### HeD—Hagerstown-Rock outcrop complex, 5 to 25 percent slopes

### **Map Unit Setting**

National map unit symbol: 15zt Elevation: 460 to 1,500 feet

Mean annual precipitation: 30 to 46 inches Mean annual air temperature: 45 to 59 degrees F

Frost-free period: 139 to 210 days

Farmland classification: Not prime farmland

### **Map Unit Composition**

Hagerstown and similar soils: 75 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Hagerstown**

### Setting

Landform: Valley floors, ridges

Landform position (two-dimensional): Shoulder, backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Linear, convex Across-slope shape: Linear, convex

Parent material: Residuum weathered from limestone

### **Typical profile**

H1 - 0 to 8 inches: silt loam H2 - 8 to 70 inches: clay

### **Properties and qualities**

Slope: 5 to 25 percent

Depth to restrictive feature: 48 to 84 inches to lithic bedrock

Drainage class: Well drained Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 2.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: High (about 10.4 inches)

#### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: B Hydric soil rating: No

### **Minor Components**

#### **Edom**

Percent of map unit: 5 percent

Landform: Hillslopes

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Convex Hydric soil rating: No

### Hublersburg

Percent of map unit: 5 percent Landform: Ridges on valleys

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear

Across-slope shape: Linear Hydric soil rating: No

### MoB-Monongahela silt loam, 2 to 10 percent slopes

### **Map Unit Setting**

National map unit symbol: 160j Elevation: 300 to 1,000 feet

Mean annual precipitation: 35 to 55 inches Mean annual air temperature: 45 to 59 degrees F

Frost-free period: 120 to 180 days

Farmland classification: All areas are prime farmland

### **Map Unit Composition**

Monongahela and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Monongahela**

### Setting

Landform: Stream terraces

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Tread

Down-slope shape: Linear Across-slope shape: Linear

Parent material: Old alluvium derived from sandstone and shale

#### Typical profile

H1 - 0 to 10 inches: silt loam
H2 - 10 to 28 inches: silt loam

H3 - 28 to 48 inches: gravelly silt loam H4 - 48 to 60 inches: gravelly silt loam

### **Properties and qualities**

Slope: 2 to 10 percent

Depth to restrictive feature: 18 to 30 inches to fragipan

Drainage class: Moderately well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to

moderately high (0.06 to 0.60 in/hr)

Depth to water table: About 18 to 36 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Low (about 5.0 inches)

#### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2e

Hydrologic Soil Group: C/D Hydric soil rating: No

### **Minor Components**

### **Tyler**

Percent of map unit: 5 percent

Hydric soil rating: No

#### Birdsboro

Percent of map unit: 5 percent Landform: Alluvial fans, terraces

Landform position (two-dimensional): Toeslope, footslope

Landform position (three-dimensional): Tread

Down-slope shape: Convex, linear Across-slope shape: Linear, convex

Hydric soil rating: No

### MuB—Murrill gravelly loam, 3 to 8 percent slopes

### **Map Unit Setting**

National map unit symbol: 160q Elevation: 200 to 1,800 feet

Mean annual precipitation: 32 to 50 inches
Mean annual air temperature: 48 to 57 degrees F

Frost-free period: 120 to 200 days

Farmland classification: All areas are prime farmland

### **Map Unit Composition**

Murrill and similar soils: 90 percent Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Murrill**

### Setting

Landform: Hills

Landform position (two-dimensional): Footslope Landform position (three-dimensional): Base slope

Down-slope shape: Linear, convex Across-slope shape: Convex, linear

Parent material: Colluvium derived from limestone, sandstone, and shale over

residuum weathered from limestone

### Typical profile

A - 0 to 9 inches: gravelly loam

Bt - 9 to 31 inches: gravelly clay loam

2Bt - 31 to 64 inches: gravelly silty clay loam

### **Properties and qualities**

Slope: 3 to 8 percent

Depth to restrictive feature: 72 to 99 inches to lithic bedrock

Drainage class: Well drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.20 to 2.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Moderate (about 6.8 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2e

Hydrologic Soil Group: B Hydric soil rating: No

### **Minor Components**

### Clarksburg

Percent of map unit: 4 percent

Landform: Valley flats

Landform position (two-dimensional): Footslope, toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave, linear Across-slope shape: Linear, concave

Hydric soil rating: No

#### **Penlaw**

Percent of map unit: 4 percent

Landform: Swales

Landform position (two-dimensional): Toeslope, footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: No

#### **Thorndale**

Percent of map unit: 2 percent

Landform: Depressions

Landform position (two-dimensional): Footslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave

Across-slope shape: Linear, concave

Hydric soil rating: Yes

### MuC—Murrill gravelly loam, 8 to 15 percent slopes

### Map Unit Setting

National map unit symbol: 160r Elevation: 200 to 1,800 feet

Mean annual precipitation: 32 to 50 inches Mean annual air temperature: 48 to 57 degrees F

Frost-free period: 120 to 200 days

Farmland classification: Farmland of statewide importance

### **Map Unit Composition**

Murrill and similar soils: 90 percent Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Murrill**

### Setting

Landform: Hills

Landform position (two-dimensional): Footslope Landform position (three-dimensional): Base slope

Down-slope shape: Linear, convex Across-slope shape: Convex, linear

Parent material: Fine-loamy colluvium derived from limestone, sandstone, and

shale over clayey residuum weathered from limestone

### **Typical profile**

Ap - 0 to 9 inches: gravelly loam

Bt - 9 to 31 inches: gravelly clay loam

2Bt - 31 to 64 inches: gravelly silty clay loam

### **Properties and qualities**

Slope: 8 to 15 percent

Depth to restrictive feature: 72 to 99 inches to lithic bedrock

Drainage class: Well drained Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.20 to 2.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Moderate (about 6.8 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3e

Hydrologic Soil Group: B Hydric soil rating: No

### **Minor Components**

#### Clarksburg

Percent of map unit: 4 percent

Landform: Valley flats

Landform position (two-dimensional): Toeslope, footslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave, linear Across-slope shape: Linear, concave

Hydric soil rating: No

#### **Penlaw**

Percent of map unit: 4 percent

Landform: Swales

Landform position (two-dimensional): Toeslope, footslope

Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: No

#### **Thorndale**

Percent of map unit: 2 percent Landform: Depressions

Landform position (two-dimensional): Footslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave

Across-slope shape: Linear, concave

Hydric soil rating: Yes

#### Ne—Newark silt loam

### **Map Unit Setting**

National map unit symbol: 160t Elevation: 320 to 1.530 feet

Mean annual precipitation: 32 to 48 inches Mean annual air temperature: 45 to 55 degrees F

Frost-free period: 120 to 205 days

Farmland classification: Farmland of statewide importance

### **Map Unit Composition**

Newark and similar soils: 80 percent Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

#### **Description of Newark**

### Setting

Landform: Flood plains, depressions

Landform position (two-dimensional): Footslope, toeslope Landform position (three-dimensional): Side slope, head slope

Down-slope shape: Linear, concave Across-slope shape: Linear, concave

Parent material: Mixed alluvium derived from limestone, sandstone, and shale

### **Typical profile**

H1 - 0 to 8 inches: silt loam H2 - 8 to 38 inches: silt loam

H3 - 38 to 60 inches: gravelly fine sandy loam

### **Properties and qualities**

Slope: 0 to 3 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat poorly drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 2.00 in/hr)

Depth to water table: About 6 to 18 inches

Frequency of flooding: NoneOccasional

Frequency of ponding: None

Available water capacity: High (about 12.0 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: B/D Hydric soil rating: No

### **Minor Components**

#### Melvin

Percent of map unit: 15 percent

Landform: Depressions
Down-slope shape: Linear
Across-slope shape: Linear
Hydric soil rating: Yes

#### **Basher**

Percent of map unit: 5 percent

Hydric soil rating: No

### Po—Philo and Basher silt loams, high bottom

### **Map Unit Setting**

National map unit symbol: 1611 Elevation: 400 to 3,000 feet

Mean annual precipitation: 32 to 55 inches Mean annual air temperature: 45 to 59 degrees F

Frost-free period: 110 to 180 days

Farmland classification: All areas are prime farmland

### **Map Unit Composition**

Philo and similar soils: 45 percent Basher and similar soils: 40 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Philo**

### Setting

Landform: Flood plains

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Mountainbase

Down-slope shape: Linear Across-slope shape: Linear

Parent material: Coarse-loamy alluvium derived from sandstone and siltstone

### **Typical profile**

H1 - 0 to 7 inches: silt loam

H2 - 7 to 40 inches: silt loam

H3 - 40 to 60 inches: stratified sandy loam to gravelly loam to loam

### **Properties and qualities**

Slope: 0 to 3 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 2.00 in/hr)

Depth to water table: About 18 to 36 inches

Frequency of flooding: RareNone Frequency of ponding: None

Available water capacity: Moderate (about 7.7 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: C Hydric soil rating: No

### **Description of Basher**

### Setting

Landform: Flood plains
Down-slope shape: Linear
Across-slope shape: Linear

Parent material: Reddish alluvium derived from sedimentary rock

### **Typical profile**

H1 - 0 to 9 inches: silt loam
H2 - 9 to 30 inches: silt loam

2H3 - 30 to 60 inches: very gravelly sandy loam

### **Properties and qualities**

Slope: 0 to 3 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high

(0.60 to 2.00 in/hr)

Depth to water table: About 18 to 24 inches

Frequency of flooding: RareNone Frequency of ponding: None

Available water capacity: Moderate (about 6.3 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 2w

Hydrologic Soil Group: B/D Hydric soil rating: No

### **Minor Components**

### **Atkins**

Percent of map unit: 10 percent Landform: Depressions

Down-slope shape: Linear

Across-slope shape: Linear Hydric soil rating: Yes

### **Tyler**

Percent of map unit: 5 percent

Hydric soil rating: No

### Pu—Purdy silt loam

### **Map Unit Setting**

National map unit symbol: 1612 Elevation: 300 to 2,000 feet

Mean annual precipitation: 35 to 50 inches
Mean annual air temperature: 48 to 59 degrees F

Frost-free period: 130 to 180 days

Farmland classification: Not prime farmland

### **Map Unit Composition**

Purdy and similar soils: 90 percent Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Purdy**

### Setting

Landform: Terraces

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Tread, talf

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Slackwater clayey alluvium derived from sedimentary rock

#### Typical profile

H1 - 0 to 9 inches: silt loam H2 - 9 to 40 inches: silty clay H3 - 40 to 60 inches: silty clay

### **Properties and qualities**

Slope: 0 to 3 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

high (0.00 to 0.20 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 0.2 mmhos/cm) Available water capacity: High (about 9.1 inches)

### Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: D Hydric soil rating: Yes

### **Minor Components**

### **Tyler**

Percent of map unit: 10 percent

Hydric soil rating: No

### W-Water

### **Map Unit Setting**

National map unit symbol: td1f

Mean annual precipitation: 36 to 50 inches

Mean annual air temperature: 46 to 59 degrees F

Frost-free period: 120 to 214 days

Farmland classification: Not prime farmland

### **Map Unit Composition**

Water: 100 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

### **Description of Water**

### Setting

Parent material: Rivers streams ponds

### Properties and qualities

Runoff class: Negligible

Frequency of ponding: Frequent

# Soil Information for All Uses

## Soil Reports

The Soil Reports section includes various formatted tabular and narrative reports (tables) containing data for each selected soil map unit and each component of each unit. No aggregation of data has occurred as is done in reports in the Soil Properties and Qualities and Suitabilities and Limitations sections.

The reports contain soil interpretive information as well as basic soil properties and qualities. A description of each report (table) is included.

## **Vegetative Productivity**

This folder contains a collection of tabular reports that present vegetative productivity data. The reports (tables) include all selected map units and components for each map unit. Vegetative productivity includes estimates of potential vegetative production for a variety of land uses, including cropland, forestland, hayland, pastureland, horticulture and rangeland. In the underlying database, some states maintain crop yield data by individual map unit component. Other states maintain the data at the map unit level. Attributes are included for both, although only one or the other is likely to contain data for any given geographic area. For other land uses, productivity data is shown only at the map unit component level. Examples include potential crop yields under irrigated and nonirrigated conditions, forest productivity, forest site index, and total rangeland production under of normal, favorable and unfavorable conditions.

## Nonirrigated Yields by Map Unit

The average yields per acre that can be expected of the principal crops under a high level of management are shown in this table. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates;

suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

If yields of irrigated crops are given, it is assumed that the irrigation system is adapted to the soils and to the crops grown, that good-quality irrigation water is uniformly applied as needed, and that tillage is kept to a minimum.

Pasture yields are expressed in terms of animal unit months. An animal unit month (AUM) is the amount of forage required by one mature cow of approximately 1,000 pounds weight, with or without a calf, for 1 month.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in the table are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local office of the Natural Resources Conservation Service or of the Cooperative Extension Service can provide information about the management and productivity of the soils for those crops.

The land capability classification of map units in the survey area is shown in this table. This classification shows, in a general way, the suitability of soils for most kinds of field crops (United States Department of Agriculture, Soil Conservation Service, 1961). Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes.

In the capability system, soils are generally grouped at three levels: capability class, subclass, and unit.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

- Class 1 soils have slight limitations that restrict their use.
- Class 2 soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.
- Class 3 soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.
- Class 4 soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.
- Class 5 soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

- Class 6 soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.
- Class 7 soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.
- Class 8 soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, e, w, s, or c, to the class numeral, for example, 2e. The letter e shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; w shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); s shows that the soil is limited mainly because it is shallow, droughty, or stony; and c, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by w, s, or c because the soils in class 5 are subject to little or no erosion.

Capability units are soil groups within a subclass. The soils in a capability unit are enough alike to be suited to the same crops and pasture plants, to require similar management, and to have similar productivity. Capability units are generally designated by adding an Arabic numeral to the subclass symbol, for example, 2e-4 and 3e-6. These units are not given in all soil surveys.

#### Reference:

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210.

### Report—Nonirrigated Yields by Map Unit

Nonirrigated Yields by Map Unit-Huntingdon County, Pennsylvania								
Map symbol and soil name	Land capability	Alfalfa hay	Corn	Corn silage	Soybeans	Wheat		
		Tons	Ви	Tons	Ви	Ви		
BuB—Buchanan gravelly loam, 3 to 8 percent slopes		3.50	100	_	_	40		
Buchanan	2e							
EgC—Edom-Weikert complex, 8 to 15 percent slopes		3.50	90	18.00	_	35		
Edom Weikert	3e 4e							
EgD—Edom-Weikert complex, 15 to 25 percent slopes		3.00	80	16.00	_	35		
Edom Weikert	6e 6e							
EgF—Edom-Weikert complex, 25 to 60 percent slopes		_	_	_	_	_		
Edom Weikert	7e 6e							
HeD—Hagerstown-Rock outcrop complex, 5 to 25 percent slopes		4.00	110	22.00	_	35		
Hagerstown	4e							
MoB—Monongahela silt loam, 2 to 10 percent slopes		3.50	100	20.00	_	40		
Monongahela	2e							
MuB—Murrill gravelly loam, 3 to 8 percent slopes		4.50	120	24.00	_	45		
Murrill	2e							
MuC—Murrill gravelly loam, 8 to 15 percent slopes		4.00	110	22.00	_	40		
Murrill	3e							
Ne—Newark silt loam		_	110	20.00	40	45		
Newark	2w							
Po—Philo and Basher silt loams, high bottom		4.50	130	_	_	45		
Philo Basher	2w 2w							
Pu—Purdy silt loam			80	16.00	_	_		
Purdy	4w							
W—Water		_	_	_	_	_		
Water	_							

# References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2\_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2\_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE\_DOCUMENTS/nrcs142p2\_052290.pdf