



United States  
Department of  
Agriculture


**NRCS**

Natural  
Resources  
Conservation  
Service

A product of the National  
Cooperative Soil Survey,  
a joint effort of the United  
States Department of  
Agriculture and other  
Federal agencies, State  
agencies including the  
Agricultural Experiment  
Stations, and local  
participants

# Custom Soil Resource Report for **Clearfield County, Pennsylvania**

## Keifer Farm Soils Report



# Preface

---

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<http://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist ([http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2\\_053951](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951)).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means

for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

# Contents

---

<b>Preface</b> .....	2
<b>Soil Map</b> .....	5
Soil Map.....	6
Legend.....	7
Map Unit Legend.....	8
Map Unit Descriptions.....	8
Clearfield County, Pennsylvania.....	10
92D—Bethesda very channery silt loam, 8 to 25 percent slopes.....	10
ErD—Ernest silt loam, 15 to 25 percent slopes.....	11
GIC—Gilpin channery silt loam, 8 to 15 percent slopes.....	12
RaB—Rayne silt loam, 3 to 8 percent slopes.....	14
RaC—Rayne silt loam, 8 to 15 percent slopes.....	15
RcD—Rayne-Gilpin complex, 15 to 25 percent slopes.....	16
<b>Soil Information for All Uses</b> .....	19
Soil Reports.....	19
Vegetative Productivity.....	19
Nonirrigated Yields by Map Unit (Soil Productivity - Keifer Farm).....	19
<b>References</b> .....	23


# Soil Map

---

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.




# Custom Soil Resource Report Soil Map




# Custom Soil Resource Report


## MAP LEGEND

### Area of Interest (AOI)

 Area of Interest (AOI)


### Soils


 Soil Map Unit Polygons


 Soil Map Unit Lines


 Soil Map Unit Points

### Special Point Features

 Blowout

 Borrow Pit


 Clay Spot


 Closed Depression

 Gravel Pit

 Gravelly Spot


 Landfill

 Lava Flow

 Marsh or swamp

 Mine or Quarry

 Miscellaneous Water


 Perennial Water

 Rock Outcrop


 Saline Spot

 Sandy Spot

 Severely Eroded Spot


 Sinkhole


 Slide or Slip


 Sodic Spot

 Spoil Area

 Stony Spot


 Very Stony Spot

 Wet Spot

 Other

 Special Line Features

### Water Features

 Streams and Canals


### Transportation

 Rails


 Interstate Highways

 US Routes

 Major Roads

 Local Roads

### Background

 Aerial Photography

## MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:20,000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service  
Web Soil Survey URL: <http://websoilsurvey.nrcs.usda.gov>  
Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Clearfield County, Pennsylvania  
Survey Area Data: Version 8, Sep 15, 2014

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Oct 6, 2011—Oct 17, 2011

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

## Map Unit Legend

Clearfield County, Pennsylvania (PA033)			
Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
92D	Bethesda very channery silt loam, 8 to 25 percent slopes	13.5	33.5%
ErD	Ernest silt loam, 15 to 25 percent slopes	0.4	1.1%
GIC	Gilpin channery silt loam, 8 to 15 percent slopes	9.3	22.9%
RaB	Rayne silt loam, 3 to 8 percent slopes	12.9	31.9%
RaC	Rayne silt loam, 8 to 15 percent slopes	0.1	0.3%
RcD	Rayne-Gilpin complex, 15 to 25 percent slopes	4.2	10.3%
<b>Totals for Area of Interest</b>		<b>40.4</b>	<b>100.0%</b>

## Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially



where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

## Clearfield County, Pennsylvania

### 92D—Bethesda very channery silt loam, 8 to 25 percent slopes

#### Map Unit Setting

*National map unit symbol:* 1lrz1  
*Elevation:* 800 to 2,800 feet  
*Mean annual precipitation:* 36 to 54 inches  
*Mean annual air temperature:* 37 to 62 degrees F  
*Frost-free period:* 130 to 160 days  
*Farmland classification:* Not prime farmland

#### Map Unit Composition

*Bethesda, unstable fill, and similar soils:* 90 percent  
*Minor components:* 10 percent  
*Estimates are based on observations, descriptions, and transects of the mapunit.*

#### Description of Bethesda, Unstable Fill

##### Setting

*Landform:* Plateaus  
*Landform position (two-dimensional):* Backslope  
*Down-slope shape:* Linear  
*Across-slope shape:* Linear  
*Parent material:* Acid loamy coal extraction mine spoil derived from interbedded sedimentary rock

##### Typical profile

*A - 0 to 7 inches:* very channery silt loam  
*C - 7 to 65 inches:* extremely channery silt loam

##### Properties and qualities

*Slope:* 8 to 25 percent  
*Depth to restrictive feature:* More than 80 inches  
*Natural drainage class:* Well drained  
*Runoff class:* Medium  
*Capacity of the most limiting layer to transmit water (Ksat):* Moderately high (0.20 to 0.60 in/hr)  
*Depth to water table:* More than 80 inches  
*Frequency of flooding:* None  
*Frequency of ponding:* None  
*Available water storage in profile:* Low (about 4.6 inches)

##### Interpretive groups

*Land capability classification (irrigated):* None specified  
*Land capability classification (nonirrigated):* 6e  
*Hydrologic Soil Group:* C

#### Minor Components

##### Gilpin

*Percent of map unit:* 5 percent  
*Landform:* Hillslopes  
*Landform position (two-dimensional):* Backslope, shoulder  
*Landform position (three-dimensional):* Side slope  
*Down-slope shape:* Convex

## Custom Soil Resource Report

*Across-slope shape:* Convex

### **Wharton**

*Percent of map unit:* 4 percent

*Landform:* Hillslopes

*Landform position (two-dimensional):* Backslope

*Landform position (three-dimensional):* Side slope

*Down-slope shape:* Linear

*Across-slope shape:* Concave

### **Wet spots**

*Percent of map unit:* 1 percent

*Landform:* Depressions

*Down-slope shape:* Concave

*Across-slope shape:* Concave

## **ErD—Ernest silt loam, 15 to 25 percent slopes**

### **Map Unit Setting**

*National map unit symbol:* l2cv

*Elevation:* 1,000 to 1,800 feet

*Mean annual precipitation:* 36 to 65 inches

*Mean annual air temperature:* 41 to 62 degrees F

*Frost-free period:* 110 to 180 days

*Farmland classification:* Not prime farmland

### **Map Unit Composition**

*Ernest and similar soils:* 90 percent

*Minor components:* 10 percent

*Estimates are based on observations, descriptions, and transects of the mapunit.*

### **Description of Ernest**

#### **Setting**

*Landform:* Hillslopes

*Landform position (two-dimensional):* Backslope

*Landform position (three-dimensional):* Side slope

*Down-slope shape:* Concave

*Across-slope shape:* Concave

*Parent material:* Acid fine-loamy colluvium derived from shale and siltstone

#### **Typical profile**

*Ap - 0 to 8 inches:* silt loam

*Bt - 8 to 24 inches:* silty clay loam

*Btx - 24 to 50 inches:* channery silt loam

*C - 50 to 74 inches:* channery silt loam

#### **Properties and qualities**

*Slope:* 15 to 25 percent

*Depth to restrictive feature:* 20 to 36 inches to fragipan

*Natural drainage class:* Moderately well drained

## Custom Soil Resource Report

*Runoff class:* High

*Capacity of the most limiting layer to transmit water (Ksat):* Moderately low to moderately high (0.06 to 0.33 in/hr)

*Depth to water table:* About 17 to 22 inches

*Frequency of flooding:* None

*Frequency of ponding:* None

*Available water storage in profile:* Low (about 3.6 inches)

### Interpretive groups

*Land capability classification (irrigated):* None specified

*Land capability classification (nonirrigated):* 4e

*Hydrologic Soil Group:* C/D

### Minor Components

#### Gilpin

*Percent of map unit:* 10 percent

*Landform:* Hills

*Landform position (two-dimensional):* Backslope

*Landform position (three-dimensional):* Side slope

*Down-slope shape:* Convex

*Across-slope shape:* Convex

## GIC—Gilpin channery silt loam, 8 to 15 percent slopes

### Map Unit Setting

*National map unit symbol:* 2snf7

*Elevation:* 810 to 3,150 feet

*Mean annual precipitation:* 38 to 50 inches

*Mean annual air temperature:* 45 to 49 degrees F

*Frost-free period:* 126 to 165 days

*Farmland classification:* Farmland of statewide importance

### Map Unit Composition

*Gilpin and similar soils:* 85 percent

*Minor components:* 15 percent

*Estimates are based on observations, descriptions, and transects of the mapunit.*

### Description of Gilpin

#### Setting

*Landform:* Hillslopes

*Landform position (two-dimensional):* Summit, shoulder

*Landform position (three-dimensional):* Interfluvium

*Down-slope shape:* Convex, linear

*Across-slope shape:* Convex

*Parent material:* Gray and brown acid fine-loamy residuum weathered from interbedded sedimentary rock

#### Typical profile

*Oi - 0 to 2 inches:* slightly decomposed plant material

## Custom Soil Resource Report

*A - 2 to 3 inches:* channery silt loam  
*E - 3 to 7 inches:* silt loam  
*Bt - 7 to 24 inches:* channery silt loam  
*C - 24 to 31 inches:* extremely channery silt loam  
*R - 31 to 41 inches:* bedrock

### Properties and qualities

*Slope:* 8 to 15 percent  
*Depth to restrictive feature:* 25 to 40 inches to lithic bedrock  
*Natural drainage class:* Well drained  
*Runoff class:* Medium  
*Capacity of the most limiting layer to transmit water (Ksat):* Moderately low to high  
(0.06 to 2.00 in/hr)  
*Depth to water table:* More than 80 inches  
*Frequency of flooding:* None  
*Frequency of ponding:* None  
*Available water storage in profile:* Low (about 4.3 inches)

### Interpretive groups

*Land capability classification (irrigated):* None specified  
*Land capability classification (nonirrigated):* 3e  
*Hydrologic Soil Group:* C

### Minor Components

#### Rayne

*Percent of map unit:* 5 percent  
*Landform:* Hillslopes  
*Landform position (two-dimensional):* Summit, shoulder  
*Landform position (three-dimensional):* Interfluve  
*Down-slope shape:* Convex, linear  
*Across-slope shape:* Convex

#### Wharton

*Percent of map unit:* 5 percent  
*Landform:* Hillslopes  
*Landform position (two-dimensional):* Summit, shoulder  
*Landform position (three-dimensional):* Interfluve  
*Down-slope shape:* Linear  
*Across-slope shape:* Concave

#### Cavode

*Percent of map unit:* 3 percent  
*Landform:* Hillslopes  
*Landform position (two-dimensional):* Summit, shoulder  
*Landform position (three-dimensional):* Interfluve  
*Down-slope shape:* Concave, linear  
*Across-slope shape:* Concave

#### Dekalb

*Percent of map unit:* 2 percent  
*Landform:* Hillslopes  
*Landform position (two-dimensional):* Summit, shoulder  
*Landform position (three-dimensional):* Interfluve  
*Down-slope shape:* Convex  
*Across-slope shape:* Convex  
*Other vegetative classification:* Very Rocky, Acid Soils (RA3)



## **RaB—Rayne silt loam, 3 to 8 percent slopes**

### **Map Unit Setting**

*National map unit symbol:* l2dj  
*Mean annual precipitation:* 37 to 65 inches  
*Mean annual air temperature:* 45 to 55 degrees F  
*Frost-free period:* 110 to 180 days  
*Farmland classification:* All areas are prime farmland

### **Map Unit Composition**

*Rayne and similar soils:* 90 percent  
*Minor components:* 10 percent  
*Estimates are based on observations, descriptions, and transects of the mapunit.*

### **Description of Rayne**

#### **Setting**

*Landform:* Hills  
*Landform position (two-dimensional):* Summit  
*Landform position (three-dimensional):* Interfluve  
*Down-slope shape:* Convex  
*Across-slope shape:* Convex  
*Parent material:* Acid loamy residuum weathered from sedimentary rock

#### **Typical profile**

*H1 - 0 to 9 inches:* silt loam  
*H2 - 9 to 38 inches:* channery silt loam  
*H3 - 38 to 60 inches:* very channery silt loam  
*H4 - 60 to 64 inches:* bedrock

#### **Properties and qualities**

*Slope:* 3 to 8 percent  
*Depth to restrictive feature:* 40 to 72 inches to lithic bedrock  
*Natural drainage class:* Well drained  
*Runoff class:* Medium  
*Capacity of the most limiting layer to transmit water (Ksat):* Moderately low to high  
(0.06 to 2.00 in/hr)  
*Depth to water table:* More than 80 inches  
*Frequency of flooding:* None  
*Frequency of ponding:* None  
*Available water storage in profile:* Moderate (about 8.3 inches)

#### **Interpretive groups**

*Land capability classification (irrigated):* None specified  
*Land capability classification (nonirrigated):* 2e  
*Hydrologic Soil Group:* B

## Minor Components

### Wharton

*Percent of map unit:* 10 percent

*Landform:* Hills

*Landform position (two-dimensional):* Backslope

*Landform position (three-dimensional):* Interfluve, side slope, head slope

*Down-slope shape:* Concave

*Across-slope shape:* Concave

## RaC—Rayne silt loam, 8 to 15 percent slopes

### Map Unit Setting

*National map unit symbol:* l2dk

*Elevation:* 1,100 to 2,500 feet

*Mean annual precipitation:* 36 to 65 inches

*Mean annual air temperature:* 45 to 55 degrees F

*Frost-free period:* 110 to 180 days

*Farmland classification:* Farmland of statewide importance

### Map Unit Composition

*Rayne and similar soils:* 90 percent

*Minor components:* 10 percent

*Estimates are based on observations, descriptions, and transects of the mapunit.*

## Description of Rayne

### Setting

*Landform:* Hills

*Landform position (two-dimensional):* Summit

*Landform position (three-dimensional):* Side slope

*Down-slope shape:* Convex

*Across-slope shape:* Convex

*Parent material:* Acid loamy residuum weathered from sedimentary rock

### Typical profile

*H1 - 0 to 9 inches:* silt loam

*H2 - 9 to 38 inches:* channery silt loam

*H3 - 38 to 60 inches:* very channery silt loam

*H4 - 60 to 64 inches:* bedrock

### Properties and qualities

*Slope:* 8 to 15 percent

*Depth to restrictive feature:* 40 to 72 inches to lithic bedrock

*Natural drainage class:* Well drained

*Runoff class:* Medium

*Capacity of the most limiting layer to transmit water (Ksat):* Moderately low to high  
(0.06 to 2.00 in/hr)

*Depth to water table:* More than 80 inches

*Frequency of flooding:* None

## Custom Soil Resource Report

*Frequency of ponding:* None

*Available water storage in profile:* Moderate (about 8.3 inches)

### Interpretive groups

*Land capability classification (irrigated):* None specified

*Land capability classification (nonirrigated):* 3e

*Hydrologic Soil Group:* B

### Minor Components

#### Wharton

*Percent of map unit:* 5 percent

*Landform:* Hills

*Landform position (two-dimensional):* Backslope

*Landform position (three-dimensional):* Interfluve, side slope, head slope

*Down-slope shape:* Concave

*Across-slope shape:* Concave

#### Hazleton

*Percent of map unit:* 5 percent

*Landform:* Hillslopes

*Landform position (two-dimensional):* Backslope

*Landform position (three-dimensional):* Head slope, side slope

*Down-slope shape:* Convex

*Across-slope shape:* Convex

## RcD—Rayne-Gilpin complex, 15 to 25 percent slopes

### Map Unit Setting

*National map unit symbol:* l2dm

*Elevation:* 800 to 2,500 feet

*Mean annual precipitation:* 36 to 65 inches

*Mean annual air temperature:* 41 to 62 degrees F

*Frost-free period:* 110 to 180 days

*Farmland classification:* Not prime farmland

### Map Unit Composition

*Rayne and similar soils:* 45 percent

*Gilpin and similar soils:* 40 percent

*Minor components:* 15 percent

*Estimates are based on observations, descriptions, and transects of the mapunit.*

### Description of Rayne

#### Setting

*Landform:* Hills

*Landform position (two-dimensional):* Backslope

*Landform position (three-dimensional):* Side slope

*Down-slope shape:* Convex

*Across-slope shape:* Convex

*Parent material:* Acid fine-loamy residuum weathered from sandstone and siltstone

## Custom Soil Resource Report

### Typical profile

*Ap* - 0 to 8 inches: channery silt loam  
*Bt* - 8 to 47 inches: channery silty clay loam  
*C* - 47 to 55 inches: channery sandy loam  
*R* - 55 to 59 inches: bedrock

### Properties and qualities

*Slope*: 15 to 25 percent  
*Depth to restrictive feature*: 40 to 72 inches to lithic bedrock  
*Natural drainage class*: Well drained  
*Runoff class*: High  
*Capacity of the most limiting layer to transmit water (Ksat)*: Moderately low to high  
(0.06 to 2.00 in/hr)  
*Depth to water table*: More than 80 inches  
*Frequency of flooding*: None  
*Frequency of ponding*: None  
*Available water storage in profile*: Moderate (about 7.6 inches)

### Interpretive groups

*Land capability classification (irrigated)*: None specified  
*Land capability classification (nonirrigated)*: 4e  
*Hydrologic Soil Group*: B

## Description of Gilpin

### Setting

*Landform*: Hills  
*Landform position (two-dimensional)*: Backslope  
*Landform position (three-dimensional)*: Side slope  
*Down-slope shape*: Convex  
*Across-slope shape*: Convex  
*Parent material*: Acid fine-loamy residuum weathered from shale and siltstone

### Typical profile

*Ap* - 0 to 6 inches: channery silt loam  
*Bt* - 6 to 24 inches: channery silt loam  
*C* - 24 to 28 inches: channery sandy loam  
*R* - 28 to 34 inches: bedrock

### Properties and qualities

*Slope*: 15 to 25 percent  
*Depth to restrictive feature*: 20 to 40 inches to lithic bedrock  
*Natural drainage class*: Well drained  
*Runoff class*: High  
*Capacity of the most limiting layer to transmit water (Ksat)*: Moderately high to high  
(0.20 to 2.00 in/hr)  
*Depth to water table*: More than 80 inches  
*Frequency of flooding*: None  
*Frequency of ponding*: None  
*Available water storage in profile*: Low (about 3.8 inches)

### Interpretive groups

*Land capability classification (irrigated)*: None specified  
*Land capability classification (nonirrigated)*: 4e  
*Hydrologic Soil Group*: C

**Minor Components**

**Wharton**

*Percent of map unit:* 15 percent

*Landform:* Hillslopes

*Landform position (two-dimensional):* Backslope

*Landform position (three-dimensional):* Side slope

*Down-slope shape:* Linear

*Across-slope shape:* Concave



# **Soil Information for All Uses**

---

## **Soil Reports**

The Soil Reports section includes various formatted tabular and narrative reports (tables) containing data for each selected soil map unit and each component of each unit. No aggregation of data has occurred as is done in reports in the Soil Properties and Qualities and Suitabilities and Limitations sections.

The reports contain soil interpretive information as well as basic soil properties and qualities. A description of each report (table) is included.

## **Vegetative Productivity**

This folder contains a collection of tabular reports that present vegetative productivity data. The reports (tables) include all selected map units and components for each map unit. Vegetative productivity includes estimates of potential vegetative production for a variety of land uses, including cropland, forestland, hayland, pastureland, horticulture and rangeland. In the underlying database, some states maintain crop yield data by individual map unit component. Other states maintain the data at the map unit level. Attributes are included for both, although only one or the other is likely to contain data for any given geographic area. For other land uses, productivity data is shown only at the map unit component level. Examples include potential crop yields under irrigated and nonirrigated conditions, forest productivity, forest site index, and total rangeland production under of normal, favorable and unfavorable conditions.

## **Nonirrigated Yields by Map Unit (Soil Productivity - Keifer Farm)**

The average yields per acre that can be expected of the principal crops under a high level of management are shown in this table. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-

## Custom Soil Resource Report

yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, barnyard manure, and green manure crops; and harvesting that ensures the smallest possible loss.

If yields of irrigated crops are given, it is assumed that the irrigation system is adapted to the soils and to the crops grown, that good-quality irrigation water is uniformly applied as needed, and that tillage is kept to a minimum.

Pasture yields are expressed in terms of animal unit months. An animal unit month (AUM) is the amount of forage required by one mature cow of approximately 1,000 pounds weight, with or without a calf, for 1 month.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in the table are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local office of the Natural Resources Conservation Service or of the Cooperative Extension Service can provide information about the management and productivity of the soils for those crops.

The land capability classification of map units in the survey area is shown in this table. This classification shows, in a general way, the suitability of soils for most kinds of field crops (United States Department of Agriculture, Soil Conservation Service, 1961). Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive landforming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes.

In the capability system, soils are generally grouped at three levels: capability class, subclass, and unit.

*Capability classes*, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

- Class 1 soils have slight limitations that restrict their use.
- Class 2 soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.
- Class 3 soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.
- Class 4 soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.
- Class 5 soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

## Custom Soil Resource Report

- Class 6 soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.
- Class 7 soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.
- Class 8 soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

*Capability subclasses* are soil groups within one class. They are designated by adding a small letter, *e*, *w*, *s*, or *c*, to the class numeral, for example, 2*e*. The letter *e* shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; *w* shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); *s* shows that the soil is limited mainly because it is shallow, droughty, or stony; and *c*, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by *w*, *s*, or *c* because the soils in class 5 are subject to little or no erosion.

*Capability units* are soil groups within a subclass. The soils in a capability unit are enough alike to be suited to the same crops and pasture plants, to require similar management, and to have similar productivity. Capability units are generally designated by adding an Arabic numeral to the subclass symbol, for example, 2*e*-4 and 3*e*-6. These units are not given in all soil surveys.

### Reference:

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210.

## **Report—Nonirrigated Yields by Map Unit (Soil Productivity - Keifer Farm)**

# Custom Soil Resource Report

Nonirrigated Yields by Map Unit—Clearfield County, Pennsylvania						
Map symbol and soil name	Land capability	Alfalfa hay	Corn	Grass-legume hay	Oats	Pasture
		<i>Tons</i>	<i>Bu</i>	<i>Tons</i>	<i>Bu</i>	<i>AUM</i>
92D—Bethesda very channery silt loam, 8 to 25 percent slopes		—	—	2.00	—	3.0
Bethesda, unstable fill	6e					
ErD—Ernest silt loam, 15 to 25 percent slopes		—	85	2.50	50	5.0
Ernest	4e					
GIC—Gilpin channery silt loam, 8 to 15 percent slopes		—	—	—	—	—
Gilpin	3e					
RaB—Rayne silt loam, 3 to 8 percent slopes		4.50	110	3.50	75	8.5
Rayne	2e					
RaC—Rayne silt loam, 8 to 15 percent slopes		4.50	100	3.50	70	8.5
Rayne	3e					
RcD—Rayne-Gilpin complex, 15 to 25 percent slopes		4.00	95	3.00	65	7.5
Rayne Gilpin	4e 4e					

# References

---

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. [http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2\\_054262](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_054262)

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. [http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2\\_053577](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577)

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. [http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2\\_053580](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580)

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. [http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2\\_053374](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053374)

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. <http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084>



## Custom Soil Resource Report

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. [http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2\\_054242](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242)

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. [http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2\\_053624](http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624)

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. [http://www.nrcs.usda.gov/Internet/FSE\\_DOCUMENTS/nrcs142p2\\_052290.pdf](http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf)